

Antibiogramme 2022 L'Hôpital Riviera-Chablais

Doubles, defined as identical microorganism with identical resistance profile isolated from the same patient within 365 days, are not included in the analysis.

Antibiotikum	Isolat		Achromobacter spp.		Acinetobacter spp.		Citrobacter freundii complex		Citrobacter koseri		Enterobacter spp.		Escherichia coli		Haemophilus influenzae		Hafnia species		Klebsiella aerogenes		Klebsiella oxytoca	
	% S	n	% S	n	% S	n	% S	n	% S	n	% S	n	% S	n	% S	n	% S	n	% S	n	% S	n
Penicillin																						
Amoxicillin					0.0%	51	1.7%	59	0.0%	136	53.2%	1858					0.0%	9	0.0%	38	0.0%	70
Ampicillin													63.9%	72								
Amoxicillin-clavulanic acid					3.9%	51	96.7%	60	0.7%	137	83.7%	1764	80.6%	72	0.0%	9	0.0%	38	92.9%	70		
Piperacillin-tazobactam	100%	7			80.4%	51	98.3%	60	79.6%	137	94.3%	1709			66.7%	9	86.8%	38	97.1%	70		
Ceftriaxon										88.9%	9	72.4%	29	100%	72							
Ceftazidime					78.4%	51	100%	60	80.3%	137	95.4%	1838			88.9%	9	89.5%	38	100%	70		
Cefepime					96.0%	50	100%	60	97.8%	136	96.9%	1821			100%	9	100%	37	100%	70		
Ertapenem					98.0%	51	100%	60	94.9%	137	99.6%	1860			88.9%	9	100%	38	100%	70		
Imipenem	100%	7	100%	18					100%	10	100%	21										
Gentamicin			100%	18	90.2%	51	100%	60	95.6%	137	90.7%	1863			100%	9	100%	38	100%	70		
Tobramycin			100%	18																		
Amikacin			94.4%	18	100%	51	100%	60	99.3%	137	99.0%	1860			100%	9	100%	38	100%	70		
Trimethoprim-sulfamethoxazole	71.4%	7	94.4%	18	90.2%	51	100%	60	93.4%	137	77.8%	1860	72.2%	72	100%	9	97.4%	38	97.1%	70		
Erythromycin																						
Ciprofloxacin			94.4%	18	86.3%	51	100%	60	93.4%	137	84.9%	1863	95.9%	73	100%	9	97.4%	38	95.7%	70		
Levofloxacin			94.4%	18																		
Nitrofurantoin					100%	51	100%	59	92.0%	137	99.1%	1860			100%	9	92.1%	38	100%	70		
Fosfomycin					92.2%	51	100%	59	54.0%	137	97.8%	1860			22.2%	9	81.6%	38	70.0%	70		
Colistin												100%	6									

According to new EUCAST recommendations %S means percentage of isolates tested susceptible (S) or susceptible with increased dosing (I) against the specific antibiotic.

Antibiogramme 2022 L'Hôpital Riviera-Chablais

Doubles, defined as identical microorganism with identical resistance profile isolated from the same patient within 365 days, are not included in the analysis.

Antibiotikum	Klebsiella pneumoniae complex		Moraxella catarrhalis		Morganella morganii		Proteus mirabilis		Proteus vulgaris		Providencia spp.		Pseudomonas aeruginosa		Salmonella spp.		Serratia marcescens		Stenotrophomonas maltophilia	
	% S	n	% S	n	% S	n	% S	n	% S	n	% S	n	% S	n	% S	n	% S	n	% S	n
Penicillin			17.6%	17																
Amoxicillin	0.0%	361			0.0%	38	68.0%	122	0.0%	29	0.0%	6			66.7%	6	0.0%	23		
Ampicillin			17.6%	17																
Amoxicillin-clavulanic acid	86.5%	348	100%	17	0.0%	38	87.3%	126	79.3%	29	0.0%	6			100%	6	0.0%	23		
Piperacillin-tazobactam	87.0%	346			97.3%	37	100%	124	100%	28	100%	6	91.8%	219	100%	6	90.9%	22		
Ceftriaxon	55.6%	9	100%	17																
Ceftazidime	91.9%	358			97.3%	37	100%	126	100%	27	100%	6	92.7%	219	100%	6	95.5%	22	77.8%	27
Cefepime	97.2%	355			100%	38	99.2%	126	96.4%	28	100%	6	91.8%	219	100%	6	95.5%	22		
Ertapenem	98.1%	361			100%	38	100%	126	100%	29	100%	6			100%	6	100%	23		
Imipenem	40.0%	5											97.3%	184						
Gentamicin	96.7%	362			97.4%	38	88.9%	126	96.6%	29	83.3%	6			0.0%	6	95.7%	23		
Tobramycin													96.3%	219						
Amikacin	99.2%	361			100%	38	100%	126	100%	29	100%	6	97.3%	219	0.0%	6	0.0%	23		
Trimethoprim-sulfamethoxazole	89.5%	361	94.1%	17	89.5%	38	66.7%	126	86.2%	29	100%	6			85.0%	20	95.7%	23	96.3%	27
Erythromycin			100%	17																
Ciprofloxacin	90.1%	362	100%	17	92.1%	38	88.1%	126	100%	29	100%	6	89.0%	219	90.0%	20	95.7%	23		
Levofloxacin													90.4%	219					92.6%	27
Nitrofurantoin	83.9%	361			0.0%	38	0.0%	126	0.0%	29	0.0%	6			100%	6	0.0%	23		
Fosfomycin	71.7%	361			0.0%	38	86.5%	126	86.2%	29	66.7%	6			100%	6	87.0%	23		
Colistin																				

According to new EUCAST recommendations %S means percentage of isolates tested susceptible (S) or susceptible with increased dosing (I) against the specific antibiotic.

Antibiogramme 2022 L'Hôpital Riviera-Chablais

Doubles, defined as identical microorganism with identical resistance profile isolated from the same patient within 365 days, are not included in the analysis.

Antibiotikum	Isolat	Enterococcus faecalis		Enterococcus faecium		Staphylococcus aureus		Staphylococcus koagulase-negativ		Streptococcus pneumoniae	
		% S	n	% S	n	% S	n	% S	n	% S	n
Penicillin										100%	23
Ampicillin		100%	201	29.4%	51					100%	11
Amoxicillin-clavulanic acid		100%	201	28.6%	49	93.2%	355	55.8%	265	100%	11
Ceftriaxon										100%	10
Imipenem						93.2%	355	55.8%	265	100%	11
Gentamicin		92.5%	201	52.9%	51	97.8%	360	66.4%	265		
Trimethoprim-sulfamethoxazole						94.7%	356	86.8%	265	96.7%	30
Clindamycin						79.4%	355	59.2%	267		
Erythromycin						77.7%	355	46.2%	264	96.7%	30
Teicoplanin		100%	201	98.0%	51	100%	92				
Vancomycin		100%	201	96.1%	51	100%	355	99.6%	267	100%	30
Ciprofloxacin						94.4%	360	64.2%	265		
Levofloxacin						94.6%	355	64.2%	265	100%	30
Fusidic acid						93.3%	359	36.2%	265		
Rifampicin						99.4%	355	97.0%	264		

According to new EUCAST recommendations %S means percentage of isolates tested susceptible (S) or susceptible with increased dosing (I) against the specific antibiotic.