

# Acute Appendicitis: Efficient Diagnosis and Management

Matthew J. Snyder, DO, Nellis Family Medicine Residency Program, Las Vegas, Nevada

Marjorie Guthrie, MD, and Stephen Cagle, MD

Saint Louis University Southwest Illinois Family Medicine Residency, Belleville, Illinois

Appendicitis is one of the most common causes of acute abdominal pain in adults and children, with a lifetime risk of 8.6% in males and 6.7% in females. It is the most common nonobstetric surgical emergency during pregnancy. Findings from the history, physical examination, and laboratory studies aid in the diagnosis of acute appendicitis. Right lower quadrant pain, abdominal rigidity, and periumbilical pain radiating to the right lower quadrant are the best signs for ruling in acute appendicitis in adults. Absent or decreased bowel sounds, a positive psoas sign, a positive obturator sign, and a positive Rovsing sign are most reliable for ruling in acute appendicitis in children. The Alvarado score, Pediatric Appendicitis Score, and Appendicitis Inflammatory Response score incorporate common clinical and laboratory findings to stratify patients as low, moderate, or high risk and can help in making a timely diagnosis. Recommended first-line imaging consists of point-of-care or formal ultrasonography. Appendectomy via open laparotomy or laparoscopy is the standard treatment for acute appendicitis. However, intravenous antibiotics may be considered first-line therapy in selected patients. Pain control with opioids, nonsteroidal anti-inflammatory drugs, and acetaminophen should be a priority and does not result in delayed or unnecessary intervention. Perforation can lead to sepsis and occurs in 17% to 32% of patients with acute appendicitis. Prolonged duration of symptoms before surgical intervention raises the risk. In moderate- to high-risk patients, surgical consultation should be accomplished quickly to reduce morbidity and mortality resulting from perforation. (*Am Fam Physician*. 2018;98(1):25-33. Copyright © 2018 American Academy of Family Physicians.)




Illustration by John Karapelou

**Appendicitis is one** of the most common causes of acute abdominal pain, with a lifetime risk of 8.6% in males and 6.7% in females.<sup>1</sup> It is the most common nonobstetric surgical emergency during pregnancy, with an incidence of 6.3 per 10,000 pregnancies during the antepartum period (compared with 9.6 per 10,000 in nonpregnant persons) and increasing to 9.9 per 10,000 postpartum.<sup>2</sup> More than 300,000 appendectomies are performed each year in the United States, and less than 10% result in the removal of a normal appendix.<sup>2-5</sup> Appendicitis is thought to be caused by luminal obstruction from various etiologies, leading to increased mucus production and bacterial overgrowth, resulting in wall tension and, eventually, necrosis and potential perforation.<sup>6</sup>

**Additional content** at <https://www.aafp.org/afp/2018/0701/p25.html>.

**CME** This clinical content conforms to AAFP criteria for continuing medical education (CME). See CME Quiz on page 18.

**Author disclosure:** No relevant financial affiliations.

## Clinical Evaluation

### SIGNS AND SYMPTOMS

Diagnosing acute appendicitis accurately and efficiently can reduce morbidity and mortality from perforation and other complications. Individual signs and symptoms are more helpful at ruling in the diagnosis than they are at ruling it out when absent. The variable location of the appendix causes variations in the clinical presentation, making diagnosis challenging, especially in pregnant women.

Table 1 presents likelihood ratios of various signs and symptoms in adults and children.<sup>7,8</sup> The signs and symptoms that best rule in acute appendicitis in adults are **right lower quadrant pain** (positive likelihood ratio [LR+] = 7.3 to 8.5), **abdominal rigidity** (LR+ = 3.8), and **radiation of periumbilical pain to the right lower quadrant** (LR+ = 3.2).<sup>7</sup> In children, however, **absent or decreased bowel sounds** (LR+ = 3.1), a **positive psoas sign** (LR+ = 3.2), a **positive obturator sign** (LR+ = 3.5), and a **positive Rovsing sign** (LR+ = 3.5) are most reliable for ruling in acute appendicitis.<sup>8</sup>

Physical examination findings specific for acute appendicitis include the psoas sign, the obturator sign, and the

## ACUTE APPENDICITIS

Rovsing sign (increased right lower quadrant pain occurring with left lower quadrant palpation). *Figures 1 and 2* illustrate how to test for the psoas and obturator signs, which significantly increase the likelihood of appendicitis when present in children.<sup>9</sup>

### CLINICAL DECISION RULES

Several clinical decision rules that incorporate findings from the patient's history, physical examination, and laboratory tests have been developed and validated in a range of populations (*Table 2*).<sup>10-12</sup> These tools typically stratify patients into low-, moderate-, and high-risk categories, and are incorporated into recommended management strategies.

The Alvarado score (<https://www.mdcalc.com/alvarado-score-acute-appendicitis>) is an eight-item, 10-point tool that is the best studied clinical decision rule in adults and children. The Pediatric Appendicitis Score (<https://www.mdcalc.com/pediatric-appendicitis-score-pas>) includes similar clinical findings in addition to a sign more relevant in children: right lower quadrant pain with coughing, hopping, or percussion. Several studies comparing the Pediatric Appendicitis Score with the Alvarado score have validated its use in children.<sup>10,12,13</sup> Likelihood ratios for cutoffs on these scores are listed in *Table 3*.<sup>10</sup>

A newer tool, the Appendicitis Inflammatory Response score (<https://www.mdcalc.com/appendicitis-inflammatory-response-air-score>), includes fewer symptoms than the Alvarado score and Pediatric Appendicitis Score, but adds an inflammatory biomarker (C-reactive protein [CRP]) and allows for different severity levels of rebound pain, leukocytosis, CRP, and polymorphonucleocytes. When the Appendicitis Inflammatory Response score was evaluated in both adults and children, the overall likelihood ratios for **high-risk**, **moderate-risk**, and low-risk groups were **13**, **1.7**, and

**0.10**, respectively.<sup>11</sup> The prevalence of appendicitis in this study was 37%, and the likelihood of appendicitis in high-risk, moderate-risk, and low-risk groups was **88%**, **50%**, and **5%**, respectively. This tool has been compared with the Alvarado score and validated as an accurate clinical decision rule.<sup>11,14</sup>

TABLE 1

### Accuracy of History and Physical Examination Findings in the Diagnosis of Acute Appendicitis

Sign/symptom	Adult <sup>7</sup>		Child <sup>8</sup>	
	Positive likelihood ratio	Negative likelihood ratio	Positive likelihood ratio	Negative likelihood ratio
Right lower quadrant pain	7.3 to 8.5*	0 to 0.28	<b>1.4</b>	NA
Rigidity	3.8	0.82	NA	NA
Migration/periumbilical pain	3.2	0.50	1.8	0.70
Pain before vomiting†	2.8	NA	NA	NA
Psoas sign	2.4	0.90	<b>3.2</b>	0.70
Fever	1.9	0.58	1.2	0.90
Guarding	1.7 to 1.8	0 to 0.54	2.1	0.47
No similar previous pain	1.5	0.32	NA	NA
Rebound tenderness	1.1 to 6.3	0 to 0.86	2.2	NA
Anorexia	1.3	0.64	1.3	0.58
Vomiting	0.92	1.1	1.3	0.65
Rectal tenderness/obstipation	0.83 to 5.3	0.36 to 1.2	2.0	0.91
Nausea	0.69 to 1.2	0.70 to 0.84	NA	NA
Obturator sign	NA	NA	<b>3.5</b>	0.73
Rovsing sign	NA	NA	<b>3.5</b>	0.72
Absent/decreased bowel sounds	NA	NA	<b>3.1</b>	0.69
Pain with hopping/coughing/percussion	NA	NA	1.6	0.52


NA = not available.

\*—Based on heterogeneous studies.

†—Based on data from one study.

Information from references 7 and 8.

FIGURE 1




The **psoas sign**. Pain on passive extension of the right thigh. Patient lies on left side. Examiner extends patient's right thigh while applying counter resistance to the right hip (asterisk).

Illustration by Floyd E. Hosmer

Reprinted with permission from Hardin DM Jr. Acute appendicitis: review and update. Am Fam Physician. 1999;60(7):2029.

FIGURE 2



The **obturator sign**. Pain on passive internal rotation of the flexed thigh. Examiner moves lower leg laterally while applying resistance to the lateral side of the knee (asterisk), resulting in internal rotation of the femur.

Illustration by Floyd E. Hosmer

Reprinted with permission from Hardin DM Jr. Acute appendicitis: review and update. Am Fam Physician. 1999;60(7):2030.

## Laboratory and Radiologic Evaluation

### LABORATORY TESTING

Individually, the white blood cell (WBC) count and inflammatory biomarkers lack accuracy for the diagnosis of acute appendicitis. However, laboratory tests are helpful when combined with signs and symptoms in clinical decision rules, or in combination with imaging studies as part of a structured evaluation. For instance, a study of 845 persons (median age = 11; prevalence of acute appendicitis = 46.5%) found that even when the WBC count was less than 10,000 per  $\mu\text{L}$  ( $10.0 \times 10^9$  per L), 20% of patients still had acute appendicitis.<sup>15</sup> However, in patients with equivocal ultrasound findings, a WBC count less than 9,000 per  $\mu\text{L}$  ( $9.0 \times 10^9$  per L) and less than 65% polymorphonuclear leukocytes increased the negative predictive value from 41.9% to 95.8% (i.e., only 4.2% had appendicitis).

The accuracy of tests such as procalcitonin, calprotectin, CRP, and the APPY1 biomarker panel (which combines values for WBC count, CRP level, and myeloid reactive protein level) in children and adults is shown in Table 4.<sup>5,16,17</sup> In children, the APPY1 has a sensitivity of 98% when used alone and 99% when combined with ultrasonography; thus, a normal test result misses only 1% to 2% of patients with appendicitis.<sup>5,16</sup> Specificity for the APPY1 panel varies from 35% to 44%, with increasing specificity as time from symptom onset increases.<sup>3</sup>

### IMAGING

Ultrasonography, computed tomography (CT), and magnetic resonance imaging are options for the evaluation of patients with suspected acute appendicitis. When selecting an imaging modality, physicians should consider the availability of experienced sonographers, potential radiation exposure, cost, length of stay in the emergency department, and diagnostic accuracy.

TABLE 2

## Diagnostic Tools for the Evaluation of Suspected Appendicitis

Alvarado score		Pediatric Appendicitis Score		Appendicitis Inflammatory Response score	
Sign/symptom	Points	Sign/symptom	Points	Sign/symptom	Points
Migration of pain	1	Migration of pain	1	Vomiting	1
Anorexia	1	Anorexia	1	Right Iliac fossa pain	1
Nausea/vomiting	1	Nausea/vomiting	1	Rebound pain, light	1
Right lower quadrant tenderness	2	Right lower quadrant tenderness	2	Rebound pain, medium	2
Rebound pain	1	Rebound pain	2	Rebound pain, strong	3
Temperature $\geq 37.3^{\circ}\text{C}$ ( $99.1^{\circ}\text{F}$ )	1	Right lower quadrant pain with coughing/hopping/percussion	2	Temperature $\geq 38.5^{\circ}\text{C}$ ( $101.3^{\circ}\text{F}$ )	1
Leukocytosis $\geq 10,000$ per $\mu\text{L}$ ( $10.0 \times 10^9$ per L)	2	Temperature $\geq 38^{\circ}\text{C}$ ( $100.4^{\circ}\text{F}$ )	1	Leukocytosis $\geq 10,000$ to $14,900$ per $\mu\text{L}$ ( $10.0$ to $14.9 \times 10^9$ per L)	1
PMN $\geq 75\%$	1	Leukocytosis $\geq 10,000$ per $\mu\text{L}$	1	Leukocytosis $\geq 15,000$ per $\mu\text{L}$ ( $15.0 \times 10^9$ per L)	2
<b>Total possible score</b>	10	PMN $\geq 75\%$	1	PMN 70% to 84%	1
< 4 : peu probable		<b>Total possible score</b>	12	PMN $\geq 85\%$	2
4-7: suspect				CRP 10 to 49 g per L	1
>7: probable				CRP $\geq 50$ g per L	2
				<b>Total possible score</b>	12

CRP = C-reactive protein; PMN = polymorphonucleocytes.

Information from references 10 through 12.

&lt; 5 : peu probable

5-8: indéterminé

&gt;8: haut risque

(Table 5).<sup>3,18-23</sup> Although CT is the most commonly used imaging study in the evaluation of suspected appendicitis (approximately 75% of cases), the National Cancer Institute, the American Academy of Pediatrics, the American College of Radiology, and other organizations recommend ultrasonography as the initial modality, especially in children and pregnant women.<sup>18,19,24,25</sup> Overweight or obese patients are

more likely to undergo CT initially because ultrasonography is more likely to be nondiagnostic in these groups.<sup>24,26</sup>

The use of clinical decision rules in conjunction with ultrasonography reduces the use of CT in the evaluation of suspected appendicitis. A prospective cohort study of 840 children with clinically suspected appendicitis (267 of whom eventually had a confirmed diagnosis) evaluated an algo-

rithm based on the Pediatric Appendicitis Score and ultrasonography.<sup>27</sup> This strategy resulted in a large decrease in CT use (75.4% to 24.2%) and a reduction in the length of emergency department stay (6.2 to 5.8 hours). Given the slightly lower sensitivity of ultrasonography for detecting acute appendicitis, there is concern for higher rates of complications or missed cases. However, a prospective observational study of 150 children (50 of whom were diagnosed with acute appendicitis via point-of-care ultrasonography) resulted in no missed cases during the three-week follow-up period among the 100 patients who did not undergo surgery.<sup>20</sup>

TABLE 3

## Accuracy of Diagnostic Tools for the Evaluation of Suspected Acute Appendicitis

Clinical decision rule	Adults		Children	
	Likelihood ratio	Probability of appendicitis (%)*	Likelihood ratio	Probability of appendicitis (%)*
<b>Alvarado score</b>				
High risk: score $\geq 7$	3.4	87	4.2	67
Moderate risk: score 4 to 6	0.42	45	0.27	12
Low risk: score $< 4$	0.03	3.7	0.02	1.9
<b>Pediatric Appendicitis Score</b>				
High risk: score $\geq 8$	NA	NA	8.1	80
Moderate risk: score 4 to 7	NA	NA	0.7	26
Low risk: score $< 4$	NA	NA	0.13	6.0

NA = not applicable.

\*—Based on pretest probability of 33% in adults and 66% in children.

Information from reference 10.

TABLE 4

## Accuracy of Laboratory Values in the Evaluation of Suspected Acute Appendicitis

Test	Population	Positive likelihood ratio	Negative likelihood ratio	Negative predictive value (prevalence of 33%)*	Negative predictive value (prevalence of 50%)*
White blood cell count <sup>16</sup>	Adults and children with suspected appendicitis (meta-analysis of 14 studies; studies with children only excluded)				
≥ 10,000 per $\mu$ L (10.0 $\times$ 10 <sup>9</sup> per L)		2.5	0.26	12%	21%
≥ 12,000 per $\mu$ L (12.0 $\times$ 10 <sup>9</sup> per L)		2.8	0.48	19%	32%
≥ 14,000 per $\mu$ L (14.0 $\times$ 10 <sup>9</sup> per L)		3.0	0.69	26%	41%
C-reactive protein level <sup>16</sup>	Adults and children with suspected appendicitis (meta-analysis of 14 studies; studies with children only excluded)				
> 10 mg per L (95.24 nmol per L)		2.0	0.32	14%	24%
> 20 mg per L (190.48 nmol per L)		2.4	0.47	19%	32%
APPY1 biomarker panel <sup>17</sup>	Adults with suspected appendicitis (n = 422)	1.5	0.07	3%	7%
APPY1 biomarker panel <sup>5</sup>	Children with suspected appendicitis (n = 185)	1.7	0.06	3%	6%
APPY1 biomarker panel plus absolute neutrophils < 7,500 per $\mu$ L (7.5 $\times$ 10 <sup>9</sup> per L) <sup>5</sup>	Children with suspected appendicitis (n = 185)	1.6	0.01	1%	1%

\*—Negative predictive value is the probability of acute appendicitis with a negative test.

Information from references 5, 16, and 17.

## Treatment

### PAIN MANAGEMENT

A meta-analysis of nine randomized controlled trials showed that the use of opioids did not significantly increase the risk of delayed or unnecessary surgery in 862 adults and children with acute abdominal pain.<sup>28</sup> Acetaminophen and nonsteroidal anti-inflammatory drugs should also be considered for pain management in patients with suspected acute appendicitis, especially in those with contraindications to opioids. A study that randomized 107 patients with acute appendicitis to narcotics plus acetaminophen vs. placebo found that pain control does not significantly increase the risk of delayed or unnecessary intervention, and does not change the Alvarado score.<sup>29</sup>

### SURGERY

Appendectomy, via open laparotomy through a limited right lower quadrant incision or via laparoscopy, is the standard treatment for acute appendicitis.<sup>1</sup> A recent meta-analysis evaluated various outcomes for open and laparoscopic appendectomies in children and adults<sup>30</sup> (eTable A). Compared with open laparotomy, laparoscopic appendectomy resulted in a lower incidence of wound infection, fewer postoperative complications, shorter length of stay, and a faster return to activity, but a longer operation time.

### ANTIBIOTIC THERAPY

Emerging evidence suggests that antibiotic therapy may be considered a first-line and possibly sole therapy in selected patients with uncomplicated appendicitis. A meta-analysis of five randomized controlled trials compared various antibiotic treatments with appendectomy in 980 adults who had uncomplicated appendicitis.<sup>31</sup>

## WHAT IS NEW ON THIS TOPIC

### Appendicitis

A meta-analysis of five randomized controlled trials found that antibiotic treatment for adults with appendicitis resulted in decreased complications, less sick leave or disability, and less need for pain medication compared with initial appendectomy. However, 40% of patients who received antibiotic therapy required appendectomy within one year.

In a study of 375 children, risk factors for appendiceal perforation included fever, vomiting, longer duration of symptoms, elevated C-reactive protein level or white blood cell count, and ultrasound findings of free abdominal fluid, visualized perforation, or a mean appendix diameter of 11 mm or more.



TABLE 5

**Comparison of Imaging Modalities in the Evaluation of Suspected Acute Appendicitis**

Imaging modality	Positive likelihood ratio	Negative likelihood ratio	Negative appendectomy prevalence (%) <sup>*</sup>	Length of emergency department stay (minutes)	Positive findings	Representative fair price <sup>†</sup>
Ultrasonography						
Point-of-care	10.4	0.4	8.1 <sup>‡</sup>	154	Increased pelvic fluid, noncompressible tubular structure > 6 mm	\$150
Experienced sonographer	36.8	0.2		141		
Novice sonographer	6.9	0.5		170		
Formal	93.8	0.4		288		
Computed tomography			4.5	487 <sup>§</sup>	Appendix diameter > 6 mm with surrounding inflammation	\$325 to \$525
Noncontrast	6.4	0.12				
Dual contrast	8.3	0				
Rectal contrast	18.6	0.07				
Magnetic resonance imaging	19.8	0.05	—	—	Increased pelvic fluid, lymphadenopathy, terminal ileum swelling	\$650

\*—Compared with prevalence of 9.8% with no imaging.

†—Fair price represents reasonable out-of-pocket costs based on price comparisons. Actual cost will vary with insurance and by region. Source: HealthCare Bluebook, <https://healthcarebluebook.com> (accessed February 24, 2018; zip code 66211).

‡—Difference between point-of-care and formal ultrasonography not available; data presented for both modalities.

§—Differences between contrast protocols not available.

Information from references 3, and 18 through 23.

Antibiotic treatment resulted in a decreased rate of complications (odds ratio = 0.54; 95% confidence interval [CI], 0.37 to 0.78), less sick leave or disability (standard mean difference = −0.19; 95% CI, −0.33 to −0.06), and less need for pain medication (standard mean difference = −1.55; 95% CI, −1.96 to −1.14). However, 40% of patients in the antibiotic group required appendectomy in the following year, compared with 8.5% of those in the appendectomy group who required a second surgery.


More recently, an open-label multicenter randomized controlled trial with 530 adults 18 to 60 years of age who had uncomplicated appendicitis reported a 73% resolution rate with ertapenem (Invanz), 1 g per day intravenously for three days, followed by a seven-day course of levofloxacin (Levaquin), 500 mg per day, plus metronidazole (Flagyl), 500 mg three times per day.<sup>4</sup> A meta-analysis identified five studies (N = 404) comparing antibiotics with surgery in children with appendicitis.<sup>32</sup> Although there were generally similar results in the studies of adults, only one of the studies of children was a randomized controlled trial. Given the risks associated with open and laparoscopic appendectomies and the high resolution rate with intravenous antibiotics, antibiotic therapy should be considered an effective treatment option for adults and children. Patient management should always be done in consultation with the surgical team in accordance with local hospital protocols and

**BEST PRACTICES IN SURGERY****Recommendations from the Choosing Wisely Campaign**

Recommendation	Sponsoring organization
Do not perform computed tomography for evaluation of suspected appendicitis in children until after ultrasonography has been considered as an option.	American College of Radiology, American College of Surgeons

**Source:** For more information on the Choosing Wisely Campaign, see <http://www.choosingwisely.org>. For supporting citations and to search Choosing Wisely recommendations relevant to primary care, see <https://www.aafp.org/afp/recommendations/search.htm>.

**FIGURE 3**



\*—Surgical consultation appropriate at any stage.

**Algorithm for evaluation of patients with suspected appendicitis in the primary care setting. (AIR = Acute Inflammatory Response.)**

Adapted with permission from Santillanes G, Simms S, Gausche-Hill M, et al. Prospective evaluation of a clinical practice guideline for diagnosis of appendicitis in children. *Acad Emerg Med.* 2012;19(8):888.

shared decision making. *Figure 3* presents an algorithm for the evaluation of patients with suspected appendicitis presenting in the primary care setting.<sup>33</sup>

## COMPLICATIONS

Perforation is the most concerning complication of acute appendicitis and may lead to abscesses, peritonitis, bowel obstruction, fertility issues, and sepsis.<sup>6,34</sup> Perforation rates among adults range from 17% to 32%,<sup>6</sup> even with increased use of imaging, and may lead to an increased length of hospital stay, extended antibiotic administration, and more severe postoperative complications. A prospective observational study showed that four of 64 children (6%) with perforated appendices were treated with antibiotics for suspected

sepsis, even after surgery.<sup>35</sup> Patient-related risk factors for perforation include older age, three or more comorbid conditions, and male sex. Time from symptom onset to diagnosis and surgery is directly associated with perforation risk.

In an observational study of 230 children with appendicitis, a delay of more than 48 hours from symptom onset to diagnosis and surgery was associated with an increase in the perforation rate compared with those in whom diagnosis and surgery occurred within 24 hours (adjusted odds ratio = 4.9 [95% CI, 1.9 to 12] vs. 3.6 [95% CI, 1.4 to 9.2]), as well as a 56% mean increase in the length of hospital stay.<sup>6</sup> Based on a study of 375 children (26% of whom had perforation), risk factors for perforation included fever, vomiting, longer duration of symptoms, elevated CRP level or

## SORT: KEY RECOMMENDATIONS FOR PRACTICE

Clinical recommendation	Evidence rating	References
The Alvarado score, Pediatric Appendicitis Score, or Appendicitis Inflammatory Response score can be used with point-of-care or formal ultrasonography and laboratory testing to help diagnose acute appendicitis and reduce the use of computed tomography.	<b>B</b>	10-12, 14, 15, 18, 27
When skilled sonographers are available, first-line imaging for patients with suspected acute appendicitis consists of point-of-care or formal ultrasonography, especially in children and pregnant women.	<b>C</b>	18, 24, 25
Opioids, nonsteroidal anti-inflammatory drugs, or acetaminophen should be provided to patients with suspected acute appendicitis.	<b>A</b>	28, 29
Open and laparoscopic appendectomies are effective surgical techniques for the treatment of acute appendicitis.	<b>A</b>	1
Intravenous antibiotics can be used as first-line therapy in children and adults with acute appendicitis.	<b>A</b>	4, 31

**A** = consistent, good-quality patient-oriented evidence; **B** = inconsistent or limited-quality patient-oriented evidence; **C** = consensus, disease-oriented evidence, usual practice, expert opinion, or case series. For information about the SORT evidence rating system, go to <https://www.aafp.org/afpsort>.

WBC count, and ultrasound findings of free abdominal fluid, visualized perforation, or a mean appendix diameter of 11 mm or more.<sup>34</sup> Surgical consultation is recommended in these patients to determine whether they are candidates for nonsurgical treatment with intravenous antibiotics.

**This article** updates previous articles on this topic by Old, et al.,<sup>36</sup> and by Hardin.<sup>9</sup>

**Data Sources:** The primary literature search was completed with Essential Evidence Plus and included searches of the Cochrane database, PubMed, and National Guideline Clearinghouse using the term acute appendicitis. In addition, a PubMed search was completed using the terms acute appendicitis, treatment, pediatric, adults, antibiotics, perforation, ultrasound, and CT. Search dates: January 16, 2017, to April 15, 2018.

**The opinions** and assertions contained herein are the private views of the authors and are not to be construed as official or as reflecting the views of Saint Louis University, the U.S. Air Force Medical Department, or the U.S. Air Force at large.

### The Authors

**MATTHEW J. SNYDER, DO, FFAFP**, is program director of the Nellis Family Medicine Residency Program, Nellis Air Force Base, Las Vegas, Nev.

**MARJORIE GUTHRIE, MD**, is program director at Saint Louis University Southwest Illinois Family Medicine Residency, Belleville.

**STEPHEN CAGLE, MD**, is an assistant clinical professor in the Department of Family and Community Medicine at Saint Louis University Southwest Illinois Family Medicine Residency.

Address correspondence to Matthew J. Snyder, DO, 4700 N. Las Vegas Blvd., Nellis AFB, NV 89191 (e-mail: [mdrnsnyder@gmail.com](mailto:mdrnsnyder@gmail.com)). Reprints are not available from the authors.

### References

1. Jaschinski T, Mosch C, Eikermann M, et al. Laparoscopic versus open appendectomy in patients with suspected appendicitis: a systematic review of meta-analyses of randomised controlled trials. *BMC Gastroenterol*. 2015;15:48.
2. Zingone F, Sultan AA, Humes DJ, West J. Risk of acute appendicitis in and around pregnancy: a population-based cohort study from England. *Ann Surg*. 2015;261(2):332-337.
3. Cuschieri J, Florence M, Flum DR, et al.; SCOAP Collaborative. Negative appendectomy and imaging accuracy in the Washington State Surgical Care and Outcomes Assessment Program. *Ann Surg*. 2008;248(4):557-563.
4. Salminen P, Paajanen H, Rautio T, et al. Antibiotic therapy vs appendectomy for treatment of uncomplicated acute appendicitis: the APPAC randomized clinical trial. *JAMA*. 2015;313(23):2340-2348.
5. Benito J, Acedo Y, Medrano L, Barcena E, Garay RP, Arri EA. Usefulness of new and traditional serum biomarkers in children with suspected appendicitis. *Am J Emerg Med*. 2016;34(5):871-876.
6. Mandeville K, Monuteaux M, Pottker T, Bulloch B. Effects of timing to diagnosis and appendectomy in pediatric appendicitis. *Pediatr Emerg Care*. 2015;31(11):753-758.
7. Wagner JM, McKinney WP, Carpenter JL. Does this patient have appendicitis? *JAMA*. 1996;276(19):1589-1594.
8. Benabbas R, Hanna M, Shah J, Sinert R. Diagnostic accuracy of history, physical examination, laboratory tests, and point-of-care ultrasound for pediatric acute appendicitis in the emergency department: a systematic review and meta-analysis. *Acad Emerg Med*. 2017;24(5):523-551.
9. Hardin DM Jr. Acute appendicitis: review and update. *Am Fam Physician*. 1999;60(7):2027-2034.
10. Ebell MH, Shinholser J. What are the most clinically useful cutoffs for the Alvarado and Pediatric Appendicitis scores? A systematic review. *Ann Emerg Med*. 2014;64(4):365-372.e2.



## ACUTE APPENDICITIS

11. Kollár D, McCartan DP, Bourke M, Cross KS, Dowdall J. Predicting acute appendicitis? A comparison of the Alvarado score, the Appendicitis Inflammatory Response score and clinical assessment [published correction appears in *World J Surg*. 2015;39(1):112]. *World J Surg*. 2015; 39(1):104-109.
12. Pogorelić Z, Rak S, Mrklić I, Jurić I. Prospective validation of Alvarado score and Pediatric Appendicitis Score for the diagnosis of acute appendicitis in children. *Pediatr Emerg Care*. 2015;31(3):164-168.
13. Khanafer I, Martin DA, Mitra TP, et al. Test characteristics of common appendicitis scores with and without laboratory investigations: a prospective observational study. *BMC Pediatr*. 2016;16(1):147.
14. Scott AJ, Mason SE, Arunakirinathan M, Reissis Y, Kinross JM, Smith JJ. Risk stratification by the Appendicitis Inflammatory Response score to guide decision-making in patients with suspected appendicitis. *Br J Surg*. 2015;102(5):563-572.
15. Anandalwar SP, Callahan MJ, Bachur RG, et al. Use of white blood cell count and polymorphonuclear leukocyte differential to improve the predictive value of ultrasound for suspected appendicitis in children. *J Am Coll Surg*. 2015;220(6):1010-1017.
16. Andersson RE. Meta-analysis of the clinical and laboratory diagnosis of appendicitis. *Br J Surg*. 2004;91(1):28-37.
17. Huckins DS, Copeland K, Self W, et al. Diagnostic performance of a biomarker panel as a negative predictor for acute appendicitis in adult ED patients with abdominal pain. *Am J Emerg Med*. 2017;35(3):418-424.
18. Doniger SJ, Kornblith A. Point-of-care ultrasound integrated into a staged diagnostic algorithm for pediatric appendicitis. *Pediatr Emerg Care*. 2018;34(2):109-115.
19. Depinet H, von Allmen D, Towbin A, Hornung R, Ho M, Alessandrini E. Risk stratification to decrease unnecessary diagnostic imaging for acute appendicitis. *Pediatrics*. 2016;138(3):e20154031.
20. Elikashvili I, Tay ET, Tsung JW. The effect of point-of-care ultrasonography on emergency department length of stay and computed tomography utilization in children with suspected appendicitis. *Acad Emerg Med*. 2014;21(2):163-170.
21. Kearney TC, Harr BC, Obszanski JR, Schaefer RF. Is contrast indicated with an abdominal CT in a patient with a suspected appendicitis? *Evidence Based Practice*. April 2015. <https://mospace.umsystem.edu/xmlui/bitstream/handle/10355/46017/AbdominalCTs.pdf?sequence=1&isAllowed=y>. Accessed February 24, 2018.
22. Kearl YL, Claudius I, Behar S, et al. Accuracy of magnetic resonance imaging and ultrasound for appendicitis in diagnostic and nondiagnostic studies. *Acad Emerg Med*. 2016;23(2):179-185.
23. Sistrom CL, McKay NL. Costs, charges, and revenues for hospital diagnostic imaging procedures: differences by modality and hospital characteristics. *J Am Coll Radiol*. 2005;2(6):511-519.
24. Kotagal M, Richards MK, Flum DR, Acierno SP, Weinsheimer RL, Goldin AB. Use and accuracy of diagnostic imaging in the evaluation of pediatric appendicitis. *J Pediatr Surg*. 2015;50(4):642-646.
25. Smith MP, Katz DS, Lalani T, et al. ACR Appropriateness Criteria® right lower quadrant pain—suspected appendicitis. *Ultrasound Q*. 2015;31(2): 85-91.
26. Schuh S, Man C, Cheng A, et al. Predictors of non-diagnostic ultrasound scanning in children with suspected appendicitis. *J Pediatr*. 2011; 158(1):112-118.
27. Shah SR, Sinclair KA, Theut SB, Johnson KM, Holcomb GW III, St Peter SD. Computed tomography utilization for the diagnosis of acute appendicitis in children decreases with a diagnostic algorithm. *Ann Surg*. 2016; 264(3):474-481.
28. Ranji SR, Goldman LE, Simel DL, Shojania KG. Do opiates affect the clinical evaluation of patients with acute abdominal pain? *JAMA*. 2006; 296(14):1764-1774.
29. Mousavi SM, Paydar S, Tahmasebi S, Ghahramani L. The effects of intravenous acetaminophen on pain and clinical findings of patients with acute appendicitis; a randomized clinical trial. *Bull Emerg Trauma*. 2014; 2(1):22-26.
30. Dai L, Shuai J. Laparoscopic versus open appendectomy in adults and children: a meta-analysis of randomized controlled trials. *United European Gastroenterol J*. 2017;5(4):542-553.
31. Mason RJ, Moazzez A, Sohn H, Katkhouda N. Meta-analysis of randomized trials comparing antibiotic therapy with appendectomy for acute uncomplicated (no abscess or phlegmon) appendicitis. *Surg Infect (Larchmt)*. 2012;13(2):74-84.
32. Huang L, Yin Y, Yang L, Wang C, Li Y, Zhou Z. Comparison of antibiotic therapy and appendectomy for acute uncomplicated appendicitis in children: a meta-analysis. *JAMA Pediatr*. 2017;171(5):426-434.
33. Santillanes G, Simms S, Gausche-Hill M, et al. Prospective evaluation of a clinical practice guideline for diagnosis of appendicitis in children. *Acad Emerg Med*. 2012;19(8):886-893.
34. van den Bogaard VA, Euser SM, van der Ploeg T, et al. Diagnosing perforated appendicitis in pediatric patients: a new model. *J Pediatr Surg*. 2016;51(3):444-448.
35. Drake FT, Mottey NE, Farrokhi ET, et al. Time to appendectomy and risk of perforation in acute appendicitis. *JAMA Surg*. 2014;149(8):837-844.
36. Old JL, Dusing RW, Yap W, Dirks J. Imaging for suspected appendicitis. *Am Fam Physician*. 2005;71(1):71-78.

## ACUTE APPENDICITIS

eTABLE A

### Outcomes of Open vs. Laparoscopic Appendectomy

Outcome	Adults (95% CI)	Children (95% CI)	Overall (95% CI)
Wound infection	OR = 0.38 (0.27 to 0.54)	NS	OR = 0.38 (0.28 to 0.53)
Intra-abdominal abscess	NS	NS	NS
Postoperative complication	OR = 0.62 (0.4 to 0.96)	NS	OR = 0.64 (0.44 to 0.93)
Reoperation	NS	NS	NS
Operation time	WMD = 10.49 (5.05 to 15.92)	WMD = 16.91 (11.96 to 21.86)	WMD = 11.59 (6.65 to 16.53)
Postoperative stay	MD = -0.78 (-1.38 to -0.17)	NS	MD = -0.79 (-1.35 to -0.23)
Return to activity	MD = -3.93 (-6.15 to -1.7)	NS	MD = -5.45 (-8.98 to -1.91)

**Note:** OR < 1 or MD < 0 favors laparoscopic appendectomy.

CI = confidence interval; MD = mean difference; NS = nonsignificant; OR = odds ratio; WMD = weighted mean difference.

Information from Dai L, Shuai J. Laparoscopic versus open appendectomy in adults and children: a meta-analysis of randomized controlled trials. United European Gastroenterol J. 2017;5(4):542-553.