

Short-course antibiotics for acute otitis media (Review)

Kozyrskyj AL, Klassen TP, Moffatt M, Harvey K



**THE COCHRANE
COLLABORATION®**

This is a reprint of a Cochrane review, prepared and maintained by The Cochrane Collaboration and published in *The Cochrane Library* 2010, Issue 9

<http://www.thecochranelibrary.com>

WILEY

TABLE OF CONTENTS

HEADER	1
ABSTRACT	1
PLAIN LANGUAGE SUMMARY	2
SUMMARY OF FINDINGS FOR THE MAIN COMPARISON	3
BACKGROUND	6
OBJECTIVES	6
METHODS	6
RESULTS	8
Figure 1.	11
Figure 2.	12
Figure 3.	13
ADDITIONAL SUMMARY OF FINDINGS	13
DISCUSSION	18
AUTHORS' CONCLUSIONS	19
ACKNOWLEDGEMENTS	19
REFERENCES	19
CHARACTERISTICS OF STUDIES	24
DATA AND ANALYSES	57
Analysis 1.1. Comparison 1 Short-acting antibiotic ≤ 48 hours in short treatment arm, Outcome 1 Treatment failure at 1 month or less.	66
Analysis 2.1. Comparison 2 Short-acting antibiotic > 48 hours short-term treatment, Outcome 1 Treatment failure at 1 month or less.	67
Analysis 2.2. Comparison 2 Short-acting antibiotic > 48 hours short-term treatment, Outcome 2 Treatment failure at 8 to 19 days.	68
Analysis 2.3. Comparison 2 Short-acting antibiotic > 48 hours short-term treatment, Outcome 3 Treatment failure at 20 to 30 days.	69
Analysis 2.4. Comparison 2 Short-acting antibiotic > 48 hours short-term treatment, Outcome 4 Treatment failure at 3 months or less.	70
Analysis 2.5. Comparison 2 Short-acting antibiotic > 48 hours short-term treatment, Outcome 5 Treatment failure at 90 days.	71
Analysis 2.6. Comparison 2 Short-acting antibiotic > 48 hours short-term treatment, Outcome 6 Treatment failure at 30 to 45 days.	72
Analysis 3.1. Comparison 3 Short-acting antibiotic > 48 hours short-term treatment, < 2 yrs old, Outcome 1 Treatment failure at 1 month or less.	73
Analysis 4.1. Comparison 4 Short-acting antibiotic > 48 hours short-term treatment, ≥ 2 yrs old, Outcome 1 Treatment failure at 1 month or less.	74
Analysis 5.1. Comparison 5 Short-acting antibiotic > 48 hours short-term treatment, perforated eardrum, Outcome 1 Treatment failure at 1 month or less.	75
Analysis 6.1. Comparison 6 Short-acting antibiotic > 48 hours short-term treatment, non-perforated eardrum, Outcome 1 Treatment failure at 1 month or less.	75
Analysis 7.1. Comparison 7 Short-acting antibiotic > 48 hours short-term treatment; Sensitivity Analysis: include chronic OM, Outcome 1 Treatment failure at 1 month or less.	76
Analysis 7.2. Comparison 7 Short-acting antibiotic > 48 hours short-term treatment; Sensitivity Analysis: include chronic OM, Outcome 2 Treatment failure at 20 to 30 days.	77
Analysis 8.1. Comparison 8 Short-acting antibiotic > 48 hours short-term treatment; Sensitivity Analysis: exclude chronic OM, Outcome 1 Treatment failure at 1 month or less.	78
Analysis 8.2. Comparison 8 Short-acting antibiotic > 48 hours short-term treatment; Sensitivity Analysis: exclude chronic OM, Outcome 2 Treatment failure at 20 to 30 days.	79
Analysis 9.1. Comparison 9 Short-acting antibiotic > 48 hours short-term treatment; Sensitivity analysis: outcome only if "cured", Outcome 1 Treatment failure at 1 month or less.	80

Analysis 9.2. Comparison 9 Short-acting antibiotic > 48 hours short-term treatment; Sensitivity analysis: outcome only if “cured”, Outcome 2 Treatment failure at 20 to 30 days.	81
Analysis 10.1. Comparison 10 Short-acting antibiotic > 48 hours short-term treatment; Sensitivity analysis: same antibiotic in treatment arms, Outcome 1 Treatment failure at 1 month or less.	82
Analysis 10.2. Comparison 10 Short-acting antibiotic > 48 hours short-term treatment; Sensitivity analysis: same antibiotic in treatment arms, Outcome 2 Treatment failure at 8 to 19 days.	83
Analysis 10.3. Comparison 10 Short-acting antibiotic > 48 hours short-term treatment; Sensitivity analysis: same antibiotic in treatment arms, Outcome 3 Treatment failure at 20 to 30 days.	84
Analysis 10.4. Comparison 10 Short-acting antibiotic > 48 hours short-term treatment; Sensitivity analysis: same antibiotic in treatment arms, Outcome 4 Treatment failure at 3 months or less.	85
Analysis 10.5. Comparison 10 Short-acting antibiotic > 48 hours short-term treatment; Sensitivity analysis: same antibiotic in treatment arms, Outcome 5 Treatment failure at 90 days.	86
Analysis 10.6. Comparison 10 Short-acting antibiotic > 48 hours short-term treatment; Sensitivity analysis: same antibiotic in treatment arms, Outcome 6 Treatment failure at 30 to 45 days.	87
Analysis 11.1. Comparison 11 Short-acting antibiotic > 48 hours short-term treatment, excluding amoxicillin-clavulanate, Outcome 1 Gastrointestinal adverse effects.	88
Analysis 12.1. Comparison 12 Short-acting antibiotic > 48 hours short-term treatment; Sensitivity analysis: treatment failure at 1 month or less and risk of bias, Outcome 1 Sensitivity analysis: allocation concealment unclear risk of bias.	89
Analysis 12.2. Comparison 12 Short-acting antibiotic > 48 hours short-term treatment; Sensitivity analysis: treatment failure at 1 month or less and risk of bias, Outcome 2 Sensitivity analysis: allocation concealment low risk of bias.	90
Analysis 12.3. Comparison 12 Short-acting antibiotic > 48 hours short-term treatment; Sensitivity analysis: treatment failure at 1 month or less and risk of bias, Outcome 3 Sensitivity analysis: blinding high and unclear risk of bias.	91
Analysis 12.4. Comparison 12 Short-acting antibiotic > 48 hours short-term treatment; Sensitivity analysis: treatment failure at 1 month or less and risk of bias, Outcome 4 Sensitivity analysis: blinding low risk of bias.	92
Analysis 13.1. Comparison 13 Short-acting antibiotic > 48 hours short-term treatment; Sensitivity analysis amoxil-clav. both arms, Outcome 1 Amoxil-clav. 5 versus 10 days, treatment failure 1 month or less.	93
Analysis 13.2. Comparison 13 Short-acting antibiotic > 48 hours short-term treatment; Sensitivity analysis amoxil-clav. both arms, Outcome 2 Excluding amoxil-clav. 5 versus 10 days, treatment failure 1 month or less.	94
Analysis 14.1. Comparison 14 Short-acting antibiotic > 48 hours short-term treatment, => 2 yrs old; Sensitivity analysis, Outcome 1 Treatment failure at 1 month or less.	95
Analysis 15.1. Comparison 15 Short-acting antibiotic > 48 hours short-term treatment, < 2 yrs old; Sensitivity analysis, Outcome 1 Treatment failure at 1 month or less.	96
Analysis 16.1. Comparison 16 Short-acting antibiotic, > 48 hours short-term treatment, adverse GI effects, Outcome 1 Gastrointestinal adverse effects.	97
Analysis 17.1. Comparison 17 Ceftriaxone, Outcome 1 Treatment failure at 1 month or less.	98
Analysis 17.2. Comparison 17 Ceftriaxone, Outcome 2 Treatment failure at 3 months or less.	99
Analysis 17.3. Comparison 17 Ceftriaxone, Outcome 3 Gastrointestinal adverse effects.	99
Analysis 18.1. Comparison 18 Azithromycin single-dose short-term treatment, Outcome 1 Treatment failure at 25 to 32 days.	100
Analysis 18.2. Comparison 18 Azithromycin single-dose short-term treatment, Outcome 2 Treatment failure at 25 to 32 days, =< 2 yrs old.	101
Analysis 18.3. Comparison 18 Azithromycin single-dose short-term treatment, Outcome 3 Treatment failure at 25 to 32 days, > 2 yrs old.	102
Analysis 18.4. Comparison 18 Azithromycin single-dose short-term treatment, Outcome 4 Gastrointestinal adverse effects.	102
Analysis 19.1. Comparison 19 Azithromycin 3 to 5 days short-term treatment, Outcome 1 Treatment failure at 1 month or less.	103
Analysis 19.2. Comparison 19 Azithromycin 3 to 5 days short-term treatment, Outcome 2 Treatment failure at 8 to 19 days.	104
Analysis 19.3. Comparison 19 Azithromycin 3 to 5 days short-term treatment, Outcome 3 Treatment failure at 20 to 30 days.	105

Analysis 20.1. Comparison 20 Azithromycin 3 to 5 days short-term treatment, < 2 years old, Outcome 1 Treatment failure at 1 month or less.	106
Analysis 21.1. Comparison 21 Azithromycin 3 to 5 days short-term treatment, => 2 years old, Outcome 1 Treatment failure at 1 month or less.	106
Analysis 22.1. Comparison 22 Azithromycin 3-5 days short-term treatment; Sensitivity analysis: include chronic OM, Outcome 1 Treatment failure at 1 month or less.	107
Analysis 22.2. Comparison 22 Azithromycin 3-5 days short-term treatment; Sensitivity analysis: include chronic OM, Outcome 2 Treatment failure at 20 to 30 days.	108
Analysis 23.1. Comparison 23 Azithromycin 3 to 5 days short-term treatment; Sensitivity analysis: exclude chronic OM, Outcome 1 Treatment failure at 1 month or less.	109
Analysis 23.2. Comparison 23 Azithromycin 3 to 5 days short-term treatment; Sensitivity analysis: exclude chronic OM, Outcome 2 Treatment failure at 20 to 30 days.	110
Analysis 24.1. Comparison 24 Azithromycin 3 to 5 days short-term treatment; Sensitivity analysis: outcome only if "cured", Outcome 1 Treatment failure at 1 month or less.	111
Analysis 24.2. Comparison 24 Azithromycin 3 to 5 days short-term treatment; Sensitivity analysis: outcome only if "cured", Outcome 2 Treatment failure at 20 to 30 days.	112
Analysis 25.1. Comparison 25 Azithromycin 3 to 5 days short-term treatment; Sensitivity analysis: similar spectrum antibiotic in treatment arms, Outcome 1 Treatment failure at 20 to 30 days.	113
Analysis 25.2. Comparison 25 Azithromycin 3 to 5 days short-term treatment; Sensitivity analysis: similar spectrum antibiotic in treatment arms, Outcome 2 Treatment failure at 1 month or less.	114
Analysis 26.1. Comparison 26 Azithromycin 3 to 5 days short-term treatment; Sensitivity analysis: risk of bias, Outcome 1 Sensitivity analysis: allocation concealment unclear or high risk of bias.	115
Analysis 26.2. Comparison 26 Azithromycin 3 to 5 days short-term treatment; Sensitivity analysis: risk of bias, Outcome 2 Sensitivity analysis: allocation concealment low risk of bias.	116
Analysis 26.3. Comparison 26 Azithromycin 3 to 5 days short-term treatment; Sensitivity analysis: risk of bias, Outcome 3 Sensitivity analysis: blinding high and unclear risk of bias.	117
Analysis 26.4. Comparison 26 Azithromycin 3 to 5 days short-term treatment; Sensitivity analysis: risk of bias, Outcome 4 Sensitivity analysis: blinding low risk of bias.	118
Analysis 27.1. Comparison 27 Azithromycin 3 to 5 days short-term treatment; Sensitivity analysis: industry funding, Outcome 1 Treatment failure 1 month or less.	119
Analysis 27.2. Comparison 27 Azithromycin 3 to 5 days short-term treatment; Sensitivity analysis: industry funding, Outcome 2 Treatment failure at 8 to 19.	120
Analysis 27.3. Comparison 27 Azithromycin 3 to 5 days short-term treatment; Sensitivity analysis: industry funding, Outcome 3 Treatment failure at 20 to 30 days.	121
Analysis 28.1. Comparison 28 Azithromycin 3 to 5 days short-term treatment; Sensitivity analysis: funding not reported, Outcome 1 Treatment failure 1 month or less.	122
Analysis 28.2. Comparison 28 Azithromycin 3 to 5 days short-term treatment; Sensitivity analysis: funding not reported, Outcome 2 Treatment failure at 8 to 19 days.	123
Analysis 28.3. Comparison 28 Azithromycin 3 to 5 days short-term treatment; Sensitivity analysis: funding not reported, Outcome 3 Treatment failure at 20 to 30 days.	124
Analysis 29.1. Comparison 29 Azithromycin 3 to 5 days short-term treatment, adverse GI effects, Outcome 1 Gastrointestinal adverse effects.	125
Analysis 30.1. Comparison 30 Methods - combined control group versus twice counting tx, Outcome 1 Twice counting - short-acting antibiotic > 48 hours short-term treatment, treatment failure at 1 month or less.	126
Analysis 30.2. Comparison 30 Methods - combined control group versus twice counting tx, Outcome 2 Combined control groups - short-acting antibiotic > 48 hours short-term treatment, treatment failure 1 month or less.	127
Analysis 30.3. Comparison 30 Methods - combined control group versus twice counting tx, Outcome 3 Twice counting - short-acting antibiotic > 48 hours short-term treatment, treatment failure at 1 to 19 days.	128
Analysis 30.4. Comparison 30 Methods - combined control group versus twice counting tx, Outcome 4 Combined control groups - short-acting antibiotic > 48 hours short-term treatment, treatment failure 1 to 19 days.	129
Analysis 30.5. Comparison 30 Methods - combined control group versus twice counting tx, Outcome 5 Twice counting - short-acting antibiotic > 48 hours short-term treatment, treatment failure 20 to 30 days.	130

Analysis 30.6. Comparison 30 Methods - combined control group versus twice counting tx, Outcome 6 Combined control groups - short-acting antibiotic > 48 hours short-term treatment, treatment failure at 20 to 30 days.	131
Analysis 30.7. Comparison 30 Methods - combined control group versus twice counting tx, Outcome 7 Twice counting - short-acting antibiotic > 48 hours short-term treatment, treatment failure at 30 to 40 days.	132
Analysis 30.8. Comparison 30 Methods - combined control group versus twice counting tx, Outcome 8 Combined control groups - short-acting antibiotic > 48 hours short-term treatment, treatment failure 30 to 40 days.	133
Analysis 30.9. Comparison 30 Methods - combined control group versus twice counting tx, Outcome 9 Twice counting - short-acting antibiotic > 48 hours short-term treatment, treatment failure at 3 months or less.	134
Analysis 30.10. Comparison 30 Methods - combined control group versus twice counting tx, Outcome 10 Combined control groups - short-acting antibiotic > 48 hours short-term treatment, treatment failure at 3 months or less.	135
APPENDICES	135
FEEDBACK	143
WHAT'S NEW	144
HISTORY	144
CONTRIBUTIONS OF AUTHORS	145
DECLARATIONS OF INTEREST	145
SOURCES OF SUPPORT	146
INDEX TERMS	146

Short-course antibiotics for acute otitis media

Anita L Kozyrskyj¹, Terry P Klassen², Michael Moffatt³, Krystal Harvey⁴

¹Department of Pediatrics, Faculty of Medicine & Dentistry, University of Alberta, Edmonton, Canada. ²Manitoba Institute of Child Health, Winnipeg, Canada. ³Department of Community Health Sciences, University of Manitoba, Winnipeg, Canada. ⁴Department of Pediatrics, University of Alberta, Edmonton, Canada

Contact address: Terry P Klassen, Manitoba Institute of Child Health, 513-715 McDermot Avenue, Winnipeg, Manitoba, MB R3E 3P4, Canada. tklassen@mich.ca.

Editorial group: Cochrane Acute Respiratory Infections Group.

Publication status and date: Stable (no update expected for reasons given in 'What's new'), published in Issue 8, 2012.

Review content assessed as up-to-date: 11 November 2009.

Citation: Kozyrskyj AL, Klassen TP, Moffatt M, Harvey K. Short-course antibiotics for acute otitis media. *Cochrane Database of Systematic Reviews* 2010, Issue 9. Art. No.: CD001095. DOI: 10.1002/14651858.CD001095.pub2.

Copyright © 2012 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

ABSTRACT

Background

Acute otitis media (AOM) is a common illness during childhood, for which antibiotics are frequently prescribed.

Objectives

To determine the effectiveness of a short course of antibiotics (less than seven days) in comparison to a long course of antibiotics (seven days or greater) for the treatment of AOM in children.

Search methods

We searched the Cochrane Central Register of Controlled Trials (CENTRAL) (*The Cochrane Library* 2009, issue 4) which contains the Acute Respiratory Infections Group's Specialised Register, MEDLINE, EMBASE, MEDLINE In-Process & Other Non-Indexed Citations, CINAHL, BIOSIS Previews, OCLC Papers First and Proceedings First, Proquest Dissertations and Theses (inception to November 2009); International Pharmaceutical Abstracts, the NLM Gateway, ClinicalTrials.gov and Current Controlled Trials (inception to August 2008).

Selection criteria

Trials were included if they met the following criteria: participants aged one month to 18 years; clinical diagnosis of ear infection; no previous antimicrobial therapy; and randomisation to treatment with less than seven days versus seven days or more of antibiotics.

Data collection and analysis

The primary outcome of treatment failure was defined as the absence of clinical resolution, relapse or recurrence of AOM during one month following initiation of therapy. Treatment outcomes were extracted from individual studies and combined in the form of a summary odds ratio (OR). A summary OR of 1.0 indicates that the treatment failure rate following less than seven days of antibiotic treatment was similar to the failure rate following seven days or more of treatment.

Main results

This update included 49 trials containing 12,045 participants. Risk of treatment failure was higher with short courses of antibiotics (OR 1.34, 95% CI 1.15 to 1.55) at one month after initiation of therapy (21% failure with short-course treatment and 18% with long-course; absolute difference of 3% between groups). There were no differences found when examining treatment with ceftriaxone for less than seven days (30% failure in those receiving ceftriaxone and 27% in short-acting antibiotics administered for seven days or more) or azithromycin for less than seven days (18% failure in both those receiving azithromycin and short-acting antibiotics administered for seven days or more) with respect to risk of treatment failure at one month or less. Significant reductions in gastrointestinal adverse events were observed for treatment with short-acting antibiotics and azithromycin.

Authors' conclusions

Clinicians need to evaluate whether the minimal short-term benefit from longer treatment of antibiotics is worth exposing children to a longer course of antibiotics.

PLAIN LANGUAGE SUMMARY

Short course antibiotics for healthy children with uncomplicated acute otitis media

Acute otitis media (AOM), or middle ear infection, is a common childhood illness, with more than half of all children having at least one infection by the time they are seven. Although otitis media often resolves without treatment, it is frequently treated with antibiotics. The length of treatment varies widely. This review of 49 trials found that treating children with a short course (less than seven days) of antibiotics, compared to treatment with a long course (seven days or greater) of antibiotics, increases the likelihood of treatment failure in the short term. No differences are seen one month later. The amount of gastrointestinal adverse events decreased with a shorter course of antibiotics.

SUMMARY OF FINDINGS FOR THE MAIN COMPARISON [\[Explanation\]](#)

Short-acting antibiotic > 48 hours short-term treatment for acute otitis media						
Patient or population: patients with acute otitis media						
Settings:						
Intervention: Short-acting antibiotic > 48 hours short-term treatment						
Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	Control	Short-acting antibiotic > 48 hours short-term treatment				
Treatment failure at 1 month or less	Study population		OR 1.34 (1.15 to 1.55)	5093 (16)		
	175 per 1000	221 per 1000 (196 to 247)				
	Medium risk population					
	157 per 1000	200 per 1000 (176 to 224)				
Treatment failure at 8 to 19 days	Study population		OR 1.37 (1.15 to 1.64)	3932 (11)		
	144 per 1000	187 per 1000 (162 to 216)				
	Medium risk population					
	117 per 1000	154 per 1000 (132 to 179)				

Treatment failure at 20 to 30 days	Study population		OR 1.16 (0.94 to 1.42)	2476 (9)
	203 per 1000	228 per 1000 (193 to 266)		
	Medium risk population			
	172 per 1000	194 per 1000 (163 to 228)		
Treatment failure at 3 months or less	Study population		OR 1.18 (0.98 to 1.41)	2068 (7)
	364 per 1000	403 per 1000 (359 to 447)		
	Medium risk population			
	364 per 1000	403 per 1000 (359 to 447)		
Treatment failure at 90 days	Study population		OR 1.16 (0.65 to 2.06)	207 (2)
	327 per 1000	360 per 1000 (240 to 500)		
	Medium risk population			
	311 per 1000	344 per 1000 (227 to 482)		
Treatment failure at 30 to 45 days	Study population		OR 1.18 (0.97 to 1.43)	1861 (5)
	368 per 1000	407 per 1000 (361 to 454)		
	Medium risk population			

	364 per 1000	403 per 1000 (357 to 450)	
<p>*The basis for the assumed risk (e.g. the median control group risk across studies) is provided in footnotes. The corresponding risk (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).</p> <p>CI: Confidence interval; OR: Odds ratio;</p>			
<p>GRADE Working Group grades of evidence</p> <p>High quality: Further research is very unlikely to change our confidence in the estimate of effect.</p> <p>Moderate quality: Further research is likely to have an important impact on our confidence in the estimate of effect and may change the estimate.</p> <p>Low quality: Further research is very likely to have an important impact on our confidence in the estimate of effect and is likely to change the estimate.</p> <p>Very low quality: We are very uncertain about the estimate.</p>			

BACKGROUND

Description of the condition

Acute otitis media (AOM) is one of the most common childhood health problems, a leading cause of visits to a physician and the most frequent reason children receive antibiotics or undergo surgery. The cost estimates for AOM in the US are between \$3 and \$5 billion per year. Most children will have at least one episode of AOM, with the peak incidence period being between 6 and 11 months (Rovers 2004).

AOM is defined as the presence of a middle ear effusion in conjunction with signs and symptoms of inflammation in the middle ear, such as otalgia, otorrhoea, fever or irritability. There is a high rate of spontaneous resolution for AOM, but if left untreated it can occasionally lead to complications, such as acute mastoiditis. Hearing loss can lead to behavioural changes and communication delay (Rovers 2004).

Description of the intervention

The standard duration of therapy in North America is 10 days (Froom 1990). However, the optimal duration of treatment is not known and varies worldwide. Half of the practitioners in Great Britain prescribe a five-day course compared to the majority of practitioners in the Netherlands who prescribe six to seven days of antibiotic therapy. Many opt for no treatment (Froom 1990).

How the intervention might work

The decline during the 1940s and 1950s in suppurative complications of AOM throughout North America and Europe has been attributed to antibiotic therapy (Berman 1995). Evidence suggests that antibiotic-treated and untreated children living in high-income countries have similar long-term outcomes (Burke 1991; Mygind 1981; van Buchem 1981). However, a meta-analysis pooling the results of four studies suggests that although the spontaneous rate of resolution was high with placebo or no drug (81%) (95% confidence interval (CI) 69% to 94%), antimicrobial therapy increased resolution by 13.7% (95% CI 8.2% to 19.2%) (Rosenfeld 1994). A meta-analysis by Del Mar documented that compared to placebo, antibiotics reduce pain at two to seven days after the start of treatment (Del Mar 1997; Sanders 2009). The absolute benefit of this finding is 5.6% fewer children experience pain at two to seven days. Long-term outcomes were similar with and without treatment.

Why it is important to do this review

Practitioners in North America may be hesitant to discontinue prescribing antibiotics, despite findings from the medical literature that support management of AOM without antibiotics. Moreover, the therapy duration for optimal outcome continues to be a question. Expert opinion has recommended five days of antimicrobial treatment for uncomplicated otitis media in children over the age of six years (Paradise 1995; Paradise 1997). The quality of scientific evidence to support a policy for shorter courses of antibiotic treatment has been assessed (Pichichero 1997), but lacks a systematic, quantitative evaluation. Concern regarding resistant bacteria from the overuse of antibiotics (Cohen 1992; Murray 1994) and poor compliance is increasing, as is the cost of health care. Therefore, it is desirable to determine the shortest duration of antibiotic treatment that would result in favorable outcomes. We felt that a systematic review would add some objectivity to this debate.

OBJECTIVES

To determine the effectiveness of a short course of antibiotics (less than seven days) in comparison to a long course (seven days or longer) of antibiotics for the treatment of children with AOM. Subgroup analyses of children less than two years old, children with a perforated ear drum, and children with a history of recurrent otitis media were pursued to address concerns that these groups may have less favorable outcomes.

METHODS

Criteria for considering studies for this review

Types of studies

For inclusion in the review, we considered randomized controlled trials (RCTs) of the empiric treatment of AOM, comparing two antibiotic regimens of different durations, as described in the title or abstract of the clinical trial. When the duration of antibiotic treatment was not specified in the title or abstract, we retrieved the trial to verify the treatment duration in both arms. We placed no language restrictions on the selection of clinical trials (Gregoire 1995).

Types of participants

We included children aged one month to 18 years, with a clinical diagnosis of AOM and no history of immediate antibiotic use, immune deficiency, chronic disease or head and neck abnormalities.

Types of interventions

We compared antibiotic therapy of a treatment arm for less than seven days (defined as the short course), with a treatment arm greater than or equal to seven days (defined as the long course). The antibiotic may be the same or different in the two treatment arms. Studies using the antibiotic ceftributen were excluded from this 2009 update due to a lack of clinical relevance as the antibiotic is not commonly used by clinicians.

Types of outcome measures

Primary outcomes

Treatment failure, which included lack of clinical resolution, relapse or recurrence of AOM during a one-month period following the initiation of therapy. Clinical resolution meant that the presenting signs or symptoms of AOM had improved or resolved.

Secondary outcomes

The cumulative number of treatment failures, relapses and recurrences reported at time of diagnosis and again at a final evaluation point between one to three months. Middle ear effusion was not classified as a treatment failure because of its documented persistence during the course of the disease, regardless of treatment. We sought data on the number of children with persistent middle ear effusion at both of the evaluation points.

Search methods for identification of studies

Electronic searches

The original search was developed and run in 1997 (see [Appendix 1](#) for details of the search). The search was then updated and run in August 2008 for the years 1997 to 2007 (see search strategies in [Appendix 2](#)), and updated again in November 2009.

In this update we searched the Cochrane Central Register of Controlled Trials (CENTRAL) (*The Cochrane Library* 2009, issue 4) which contains the Cochrane Acute Respiratory Infections Group's Specialised Register; MEDLINE (1966 to November Week 1, 2009); EMBASE (1974 to November 2009); MEDLINE In-Process & Other Non-Indexed Citations (1966 to Week 1, 2009); International Pharmaceutical Abstracts (1970 to August Week 1, 2008); BIOSIS Previews (1969 to November 2009); CINAHL (1981 to November 2009); the NLM Gateway (1998 to August 2008); OCLC Papers First and Proceedings First (1997 to November 2009); ClinicalTrials.gov (1998 to August 2008); Proquest Dissertations and Theses (1861 to November 2009); and Current Controlled Trials (1997 to August 2008). We searched the following databases without any date restrictions in September 2007: the National Research Register; CRISP; the TRIP Database;

Scirus; and Google Scholar. We imposed no language or publication restrictions.

We ran the following search strategy in MEDLINE and CENTRAL. The MEDLINE search was combined with the Cochrane Highly Sensitive Search Strategy for identifying randomized trials in MEDLINE: sensitivity- and precision-maximising version (2008 revision); Ovid version ([Lefebvre 2008](#)). The search strategy also incorporated the child search strategy developed by Boluyt ([Boluyt 2008](#)). We adapted this search strategy to search the other databases.

MEDLINE (Ovid)

```
1 exp Otitis Media/
2 otitis media.tw.
3 (OM or OME or AOM).tw.
4 (otorrhea* or otorrhoea*).tw.
5 (ear* adj3 (infect* or acute*)).tw.
6 or/1-5
7 exp treatment outcome/
8 exp Anti-Infective Agents/
9 (antibacter* or anti-bacter* or anti bacter* or antibiotic* or anti-biotic* or anti biotic* or bacteriocid* or antimicrob* or anti-microb* or anti microb*).tw.
10 (amoxicillin* or amoxycillin* or penicillin* or cefprozil* or clarithromycin* or cefpodoxime* or cefaclor* or ceftriaxone* or azthromycin* cefixime*).tw.
11 or/7-10
12 6 and 11
```

Searching other resources

We retrieved clinical trial references and published reviews on the treatment of AOM in order to identify further trials. We then contacted primary authors for information on such trials.

Data collection and analysis

Selection of studies

In the first publication of this review ([Kozyrskyj 2000](#)) clinical trials examining the antibiotic treatment of AOM were retrieved from the medical literature search and evaluated by the review authors according to the selection criteria previously described. Decisions to include the trial were made independently. Interreviewer agreement was assessed by the kappa statistic using PC Agree Software (Stephen Walter, McMaster University, Hamilton, Ontario) and consensus was reached regarding trial inclusion.

In this updated review, two review authors (KH, NH) evaluated for inclusion in the review the clinical trials retrieved during the search update. A third review author (TK) was consulted when consensus could not be reached.

Data extraction and management

At least two review authors (KH, LB) independently performed data extraction, followed by a consensus conference to resolve any differences. We developed a standardized form to facilitate data extraction. Data extracted from each trial included:

- the number of children with no clinical resolution of AOM;
- relapse or recurrence of AOM;
- the number of children with persistent middle ear effusion; and
- the number of withdrawals from each arm of the trial.

We recorded the data as follows:

Primary outcome: evaluation points included time points until one month after initiation of therapy.

Secondary outcome: evaluation point between one to three months after initiation of therapy.

Important subgroups defined a priori for which data were sought are:

1. antibiotic use for two days or less;
2. children aged less than two years; and
3. perforated tympanic membrane ([Appelman 1991](#); [Hendrickse 1988](#); [Kaleida 1991](#); [van Buchem 1981](#)).

Descriptive data collected included treatment site, patient baseline characteristics, co-interventions, and inclusion, exclusion and outcome criteria to summarise the generalisability of included studies and to facilitate sensitivity analyses. We also recorded drug dose, route and treatment duration.

Change in statistical methods from original version of review

Two studies ([Gooch 1996](#); [Hoberman 1997](#)) in the current review contained two arms comparing long-course versus short-course antibiotics. In the first version of this review ([Kozyrskyj 2000](#)) the short-course arm was counted twice against each long-course arm of that study. In this review update, we have modified our methods and combined both long-course arms into a single comparison against the short-course arm, as recommended in the *Cochrane Handbook for Systematic Reviews of Interventions* ([Higgins 2008](#)). This prevents overweighing the short-course arm of these trials in the meta-analysis.

Assessment of risk of bias in included studies

We used the Cochrane Collaboration's 'Risk of bias' tool ([Higgins 2008](#)) to assess for bias in this updated review. Two review authors (KH, MO) independently evaluated the risk of bias of included trials and then came to a consensus. When consensus could not be reached a third review author was consulted (TK).

Measures of treatment effect

We compared the treatment failure rate following a short course of antibiotics to the treatment failure rate following a long course of antibiotics using odds ratios (ORs) for individual trial outcomes.

We determined a summary OR with 95% CI for trials pooled by antibiotic type using the Peto fixed-effect model.

Unit of analysis issues

We performed analysis on trials grouped by the pharmacokinetic behaviour of the antibiotic used in the short-course treatment arm, as follows.

1. Short-acting oral antibiotics, for example, penicillin, amoxicillin, cefaclor and cefuroxime.
2. Oral azithromycin.
3. Intramuscular ceftriaxone.

We also conducted additional meta-analyses in the short-acting antibiotic group for treatment duration less than 48 hours and more than 48 hours.

Assessment of heterogeneity

We assessed statistical heterogeneity ([Higgins 2008](#); [Laupacis 1988](#)). When significant heterogeneity existed, we examined trials for specific potential clinical differences ([Thompson 1994](#)). We also calculated summary ORs using the DerSimonian and Laird random-effects model ([DerSimonian 1986](#)). We determined the summary risk difference (difference in the failure rate between the short and long-course antibiotics) and a 95% CI for pooled group data.

Sensitivity analysis

We performed a sensitivity analyses to assess whether the results of the primary outcome for all antibiotics was sensitive to blinding, allocation concealment and publication bias. Additionally a sensitivity analysis was undertaken to assess those trials that used the same antibiotic in both the short and long courses of the trial.

RESULTS

Description of studies

See: [Characteristics of included studies](#); [Characteristics of excluded studies](#).

See [Characteristics of included studies](#) and [Characteristics of excluded studies](#) tables.

Results of the search

Updated electronic searches and handsearches from January 2000 through to November 2009 retrieved 4175 additional citations. In this updated review, we included 49 studies containing 12,045 participants. Refer to the [Characteristics of included studies](#) for detailed information on individual trials.

Included studies

We included 22 new trials, resulting in 49 eligible trials for this updated review (Adam 2000; Al-Ghamdi 1999; Arguedas 2005; Arrieta 2003; Block 2000; Block 2003; Block 2004; Catania 2004; Cohen 1998; Cohen 1999; Cohen 2000; Dagan 2000a; Dagan 2000b; de Jose 1998; Dunne 2003; Guven 2006; Hoberman 2005; Kara 1998; Oguz 2003; Pessey 1999; Varsano 1997; Wang 2004). Forty-eight of the included studies were classified as randomized controlled trials (RCTs). Randomisation was not mentioned in one study (Mohs 1993). Included sample sizes ranged from 17 participants (Puczynski 1987) to 868 participants (Hoberman 1997). The median sample size was 215 participants (Dagan 2000b). Studies were primarily conducted in North America, South America, Europe and the Middle East (Israel and Turkey). Single trials were carried out in both Egypt (Mohs 1993) and China (Wang 2004). All trials enrolled children; 39 trials enrolled children less than one year old; eight trials began enrolment at two years of age; and two trials evaluated children three years of age or less. The maximum age of enrolment was 14 years (Kafetzis 1997).

The following antibiotics were evaluated in the included studies (number of studies in brackets): amoxicillin or amoxicillin/clavulanate (32); azithromycin (21); cefaclor (8); cefdinir (2); cefixime (2); cefpodoxime (4); cefprozil (2); cefuroxime (3); ceftriaxone (9); clarithromycin (1); penicillin (3); and trimethoprim-sulfamethoxazole (1). Short-course interventions were primarily a three to five-day regimen. Two studies compared a single dose of azithromycin (Arguedas 2005; Block 2003) and two studies compared a short course of the antibiotic regimen at less than or equal to 48 hours (Meistrup-Larsen 1983; Puczynski 1987).

Short regimens of ceftriaxone or azithromycin were analyzed separately from all other antibiotics. Twenty trials employed tympanocentesis. Of these, four trials (Hendrickse 1988; Pessey 1999; Ploussard 1984; Puczynski 1987) studied short regimens of antibiotics and 15 trials evaluated azithromycin (Arguedas 1996; Arguedas 1997; Arguedas 2005; Aronovitz 1996; Arrieta 2003; Dagan 2000a; Dagan 2000b; Daniel 1993; de Jose 1998; Guven 2006; Hoberman 2005; Khurana 1996; Oguz 2003; Petalozza 1992; Principi 1995).

For the majority of studies, clinical cure or improvement at the end of treatment and at a follow-up period of 10 days to one month after beginning the study were primary outcomes. Please refer to the [Characteristics of included studies](#) table for details on individual study outcomes.

Excluded studies

We excluded two short-acting antibiotic trials because outcomes were not reported as treatment failures (Bain 1985; Jones 1986). One study (Jones 1986) reported the number of days symptoms were recorded in diaries and the percentage resolution of ear signs, and the other study (Bain 1985) reported eardrum signs and symptoms. Neither of these studies presented data that could contribute

to treatment failure as defined in this review. We excluded an additional two studies because the antibiotic used, ceftibuten, was deemed to be not clinically relevant due to its limited use in clinical practice (Roos 2000; Simon 1997), and excluded one study due to the duration of the antibiotic treatment being the same in both arms (Suzuki 2009).

Risk of bias in included studies

A detailed summary of the risk of bias assessment for each study is available in the [Characteristics of included studies](#) table.

Allocation

Adequate sequence generation was performed in 12 studies and was unclear in 37. Allocation concealment was adequate in six studies, inadequate in one study and unclear in 42 studies.

Blinding

Appropriate blinding was employed in 15 studies, there was no blinding in 24 studies and unclear blinding in 10 studies. Allocation concealment was also judged as adequate in only one of the appropriately blinded studies (Green 1993).

Incomplete outcome data

In 33 studies incomplete outcome data were appropriately addressed. Six studies did not address the incomplete outcome data appropriately and in 10 studies it was unclear if the data were appropriately reported.

Selective reporting

Thirty-eight included studies were judged free from a risk of selective reporting bias; five studies had a high risk of selective reporting bias; and six studies were unclear in risk of selective reporting bias.

Other potential sources of bias

Fifteen studies were judged as low risk for other sources of bias. In 31 studies the risk of other sources of bias was unclear, and in three studies the risk was judged as high. Unreported sources of funding were the primary reason for an unclear risk of bias.

Effects of interventions

See: [Summary of findings for the main comparison](#) Short-acting antibiotic > 48 hours short-term treatment for acute otitis media; [Summary of findings 2](#) Ceftriaxone for acute otitis media; [Summary of findings 3](#) Azithromycin 3 to 5 days short-term treatment for acute otitis media

Treatment failure

Treatment failure has been previously defined as lack of clinical resolution, relapse or recurrence of AOM during a one-month period following the initiation of therapy. See summary of findings tables for each intervention ([Summary of findings for the main comparison](#); [Summary of findings 2](#); [Summary of findings 3](#)).

Outcome of short-acting antibiotics in the short course (less than seven days)

Two trials were included in the meta-analysis for short-acting antibiotics administered for a period of less than 48 hours, and 17 trials were included in the meta-analysis of short-acting antibiotics administered for more than 48 hours.

i) Short-acting antibiotics given for less than 48 hours

A summary odds ratio (OR) of 2.99 (95% CI 1.04 to 8.54) was found for treatment failures at less than one month ([Analysis 1.1](#)). Two trials evaluating a total of 118 children were included in this meta-analysis ([Meistrup-Larsen 1983](#); [Puczynski 1987](#)). The trial by Puczynski was prematurely terminated ([Puczynski 1987](#)).

ii) Short-acting antibiotics given for more than 48 hours

Risk of treatment failure, defined as lack of clinical resolution, relapse or recurrence of AOM at one month or less, was higher with short courses of antibiotics (OR 1.34, 95% CI 1.15 to 1.55) ([Analysis 2.1](#)). Significant results were found when treatment failure was compared at a follow-up period of 19 days or less (OR 1.37, 95% CI 1.15 to 1.64) ([Analysis 2.2](#)). At 21 days there was no difference in the risk of treatment failure between short and long-course antibiotics ([Analysis 2.3](#)). At one month or less, 22 children needed treatment with short regimens of antibiotics to cause an additional treatment failure compared to longer regimens of antibiotics.

Subgroup analysis examining treatment failure in children less than two years and two years of age or older was performed. Six trials containing 614 children found non-significant results for treatment failure at one month or less in short versus long-course regimens of antibiotics ([Analysis 3.1](#)). Significant differences in outcomes between five-day and 10-day treatment among children less than two years old were reported in one trial, but no extractable subgroup data were presented ([Hoberman 1997](#)). Non-significant results were repeated in seven trials containing 1183 participants when examining children two years of age or older ([Analysis 4.1](#)). Hendrickse reported outcome data for perforated and non-perforated eardrums ([Hendrickse 1988](#)). The OR for treatment failure in children with perforated eardrums ($n = 27$) was 3.62 (95% CI 0.81 to 16.06) ([Analysis 5.1](#)) and 1.06 (95% CI 0.40 to 2.75) ([Analysis 6.1](#)) in children with non-perforated eardrums ($n = 101$).

Outcome of ceftriaxone in the short course (less than seven days)

Eight trials ($n = 1709$) were included in the meta-analysis for treatment failure at one month or less. No significant results were found (OR 1.07, 95% CI 0.86 to 1.33) ([Analysis 17.1](#)), and this lack of significance was repeated when treatment failure at three months or less was examined ([Analysis 17.2](#)).

Outcome of azithromycin in the short course (less than seven days)

A comparison of regimens of short-course azithromycin for treatment failure at one month or less yielded an OR of 1.02 (95% CI 0.87 to 1.20). This meta-analysis included 19 trials and 4354 children ([Analysis 19.1](#)). Examination of a follow-up period of 8 to 19 days (18 trials, $n = 4347$) found a significant OR of 1.27 (95% CI 1.04 to 1.55) ([Analysis 19.2](#)). Forty-four children needed treatment with short-course azithromycin so that one child may experience an additional treatment failure compared to children treated with longer courses of antibiotics. This significant result was not repeated when 20 to 30-day follow-up periods were examined ([Analysis 19.3](#)).

The odds of treatment failure with azithromycin was 1.92 (95% CI 0.73 to 5.04) ([Analysis 20.1](#)) in children less than two years ($n = 138$, 17.4%) and 1.34 (95% CI 0.61 to 2.94) ([Analysis 21.1](#)) in older children ($n = 656$) ([Principi 1995](#); [Schaad 1993](#)).

Sensitivity analysis

Assessment of risk of bias

A sensitivity analysis of allocation concealment and blinding from the risk of bias assessment and treatment failure at one month or less was completed. An OR of 1.30 (95% CI 1.09 to 1.54) was found in short-acting antibiotics for treatment failure in 14 studies ([Analysis 12.1](#)) where the risk of bias for allocation concealment was unclear and 1.45 (95% CI 1.08 to 1.93) ([Analysis 12.2](#)) in three studies where the risk of bias was low for allocation concealment. Thirteen studies had a high or unclear risk of bias in blinding and an OR of 1.18 (95% CI 1.00 to 1.40) ([Analysis 12.3](#)) compared to four studies with a low risk of bias and an OR of 2.03 (95% CI 1.48 to 2.77) ([Analysis 12.4](#)).

Due to a lack of difference between the risk of bias assessment for studies examining ceftriaxone, a sensitivity analysis was not completed for this outcome.

Seventeen studies using azithromycin as the antibiotic in the short course were judged to have a high or unclear risk of bias for allocation concealment. The OR was 1.04 (95% CI 0.88 to 1.22) ([Analysis 26.1](#)). Two studies had a low risk of bias for allocation concealment and had an OR of 0.13 (95% CI 0.02 to 0.90) ([Analysis 26.2](#)). The large difference between these two values

should be interpreted with caution given the low number of studies included in the meta-analysis of low risk of bias for allocation concealment. For blinding, 15 studies had a high or unclear risk of bias and an OR of 1.15 (95% CI 0.94 to 1.40) (Analysis 26.3) and four studies with a low risk of bias for blinding and an OR of 0.80 (95% CI 0.60 to 1.07) (Analysis 26.4).

Use of same antibiotic in both arms

A total of 10 studies utilised the same antibiotic in both the short and long courses of therapy. Nine studies including 3321 participants presented data for treatment failure at one month or less; this analysis resulted in an OR of 1.65 (95% CI 1.35 to 2.01) (Analysis 10.1). A sensitivity analysis was also undertaken to compare the two trials that used the same antibiotic (amoxicillin-clavulanate) in both arms. This analysis resulted in an OR of 1.99 (95% CI 1.44 to 2.74) (Analysis 13.1) compared to an OR of 1.20 (95%

CI 1.02 to 1.42) when excluding the above trials from the analysis (Analysis 13.2). The effect was larger with amoxicillin-clavulanate, however caution must be used when interpreting these results due to the number of subgroup analyses.

Publication bias

We produced funnel plots for the antibiotic comparisons, short-acting antibiotics (Figure 1), ceftriaxone (Figure 2) and azithromycin (Figure 3). Upon visual inspection, funnel plots suggested publication bias for the azithromycin comparison but not for the other antibiotic groups. Additional statistical tests (Duval 2000; Egger 1997) of significance did not reveal publication bias in any of the groups. A sensitivity analysis of studies declaring industry funding as compared to those not reporting a funding source was not significant (Analysis 27.1 and Analysis 28.1).

Figure 1. Funnel plot of comparison: 2 Short-acting antibiotic > 48 hours short-term treatment, outcome: 2.1 Treatment failure at 1 month or less.




Figure 2. Funnel plot of comparison: 21 Ceftriaxone, outcome: 21.1 Treatment failure at 1 month or less.





Figure 3. Funnel plot of comparison: 23 Azithromycin 3 to 5 days short-term treatment, outcome: 23.1 Treatment failure at 1 month or less.



Gastrointestinal adverse events

Outcome of short-acting antibiotics in the short course (less than seven days)

In 13 trials containing 4918 children a significant reduction in gastrointestinal adverse events was observed ([Analysis 16.1](#)). The OR was found to be 0.72 (95% CI 0.60 to 0.87) which translates to one less child experiencing an adverse event for every 29 children given a short regimen of antibiotics. A sensitivity analysis of the five trials that did not use amoxicillin-clavulanate provided non-significant results ([Analysis 11.1](#)).

Outcome of ceftriaxone in the short course (less than seven days)

One trial was included in the analysis of ceftriaxone adverse events ([Barnett 1997](#)). In this study ($n = 402$) more children experienced adverse events in the short course ceftriaxone group (OR 2.89, 95% CI 1.70 to 4.91). This OR translates to an additional child experiencing an adverse gastrointestinal event for every seven to eight children treated compared to long courses of antibiotics ([Analysis 17.3](#)).

Outcome of azithromycin in the short course (less than seven days)

In 14 trials containing 3722 children, significantly fewer gastrointestinal events were experienced in the short course of azithromycin comparison group ([Analysis 29.1](#)). The OR was 0.36 (95% CI 0.28 to 0.46) which translates to one less child experiencing an adverse event for every 14 children treated compared to longer courses of antibiotics.

ADDITIONAL SUMMARY OF FINDINGS *[Explanation]*

Ceftriaxone for acute otitis media						
Patient or population: patients with acute otitis media						
Settings:						
Intervention: Ceftriaxone						
Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	Control	Ceftriaxone				
Treatment failure at 1 month or less	Study population		OR 1.07 (0.86 to 1.33)	1709 (8)		
	270 per 1000	284 per 1000 (241 to 330)				
	Medium risk population					
	254 per 1000	267 per 1000 (226 to 312)				
Treatment failure at 3 months or less	Study population		OR 0.89 (0.66 to 1.21)	701 (3)		
	402 per 1000	374 per 1000 (307 to 449)				
	Medium risk population					
	379 per 1000	352 per 1000 (287 to 425)				
*The basis for the assumed risk (e.g. the median control group risk across studies) is provided in footnotes. The corresponding risk (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).						
CI: Confidence interval; OR: Odds ratio;						

GRADE Working Group grades of evidence

High quality: Further research is very unlikely to change our confidence in the estimate of effect.

Moderate quality: Further research is likely to have an important impact on our confidence in the estimate of effect and may change the estimate.

Low quality: Further research is very likely to have an important impact on our confidence in the estimate of effect and is likely to change the estimate.

Very low quality: We are very uncertain about the estimate.

Azithromycin 3 to 5 days short-term treatment for acute otitis media						
Patient or population: patients with acute otitis media Settings: Intervention: Azithromycin 3 to 5 days short-term treatment						
Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	Control	Azithromycin 3 to 5 days short-term treatment				
Treatment failure at 1 month or less	Study population		OR 1.02 (0.87 to 1.2)	4354 (19)		
	185 per 1000	188 per 1000 (165 to 214)				
	Medium risk population					
	61 per 1000	62 per 1000 (53 to 72)				
Treatment failure at 8 to 19 days	Study population		OR 1.27 (1.04 to 1.55)	4347 (18)		
	95 per 1000	118 per 1000 (98 to 140)				
	Medium risk population					
	56 per 1000	70 per 1000 (58 to 84)				
Treatment failure at 20 to 30 days	Study population		OR 0.98 (0.82 to 1.17)	2708 (11)		

	265 per 1000	261 per 1000 (228 to 297)
	Medium risk population	
	273 per 1000	269 per 1000 (235 to 305)

*The basis for the **assumed risk** (e.g. the median control group risk across studies) is provided in footnotes. The **corresponding risk** (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

CI: Confidence interval; **OR:** Odds ratio;

GRADE Working Group grades of evidence

High quality: Further research is very unlikely to change our confidence in the estimate of effect.

Moderate quality: Further research is likely to have an important impact on our confidence in the estimate of effect and may change the estimate.

Low quality: Further research is very likely to have an important impact on our confidence in the estimate of effect and is likely to change the estimate.

Very low quality: We are very uncertain about the estimate.

DISCUSSION

Summary of main results

A reduction in the treatment of AOM from 10 to five days of short-acting antibiotics may slightly increase the risk of a child experiencing signs and symptoms, relapse or re-infection at eight to 19 days (OR 1.37, 95% CI 1.15 to 1.64). By 30 days following initiation of therapy, a longer course of short-acting antibiotics is comparable to a five-day course in terms of these outcomes (OR 1.17, 95% CI 0.95 to 1.43). When studies comparing the same antibiotics in both arms were analyzed the results favoured the use of longer regimens of therapy (OR 1.65, 95% CI 1.35 to 2.01). Notably 22 studies published after 1998 were added to the updated meta-analysis and brought the 30-day findings closer to equivalence. Studies that had low risk of bias for the component of blinding had significantly higher risks of treatment failure (OR 2.03, 95% CI 1.48 to 2.77) compared to those with high risk of bias for blinding (OR 1.19, 95% CI 1.01 to 1.41). The long-term comparability between a short and long course of antibiotics is biologically plausible, on the basis of: 1) spontaneous resolution of untreated AOM (Del Mar 1997; Rosenfeld 1994); 2) early eradication of pathogens after three to five days of treatment (Howie 1969); 3) poorer penetration of the antibiotic into the ear with continued administration as inflammation decreases (Canafax 1991); and 4) treatment of children without AOM because of diagnostic uncertainty (Froom 1990).

Overall completeness and applicability of evidence

Appreciating that a shortened course of antibiotics may protect the child from developing resistant microorganisms (Kozyrskyj 1998), we report these results 12 years later when many clinical practice guidelines are proposing no antibiotic treatment or 'watchful waiting' as a first-line approach to treating AOM. The American Academy of Pediatrics recommends reserving antibiotic therapy for children that are less than or equal to two years with severe disease (Pediatric Guidelines 2004). More recently the Canadian Paediatric Society recommended a 'wait and watch' approach for children over six months with uncomplicated, non-severe disease (Forgie 2009). This is consistent with a Cochrane Review examining the effectiveness of antibiotics for AOM (Sanders 2009) and an individual patient meta-analysis (Koopman 2008; Spiro 2008; Vouloumanou 2009). If antibiotics are used, our results indicate that a long course of treatment can minimise the risks of treatment failure or recurrence post-treatment, but may not make a difference in the long term.

This updated review also found no significant differences when examining short courses of ceftriaxone. However, in the one study where adverse effects were examined, they were much higher in the ceftriaxone group. As administering ceftriaxone also requires an

injection, this does not seem to be a good option for management of AOM.

Quality of the evidence

Higher risk of treatment failure occurred with three to five days of azithromycin (OR 1.27, 95% CI 1.04 to 1.55) at 8 to 19 days after the initiation of treatment. This significant result was not repeated when examining treatment failure at 20 to 30 days (OR 0.98, 95% CI 0.82 to 1.17). Outcome differences noted between the evaluations at eight to 19 days and 20 to 30 days likely reflects bias in the timing of evaluation. Children treated with a long course of antibiotics had fewer days to experience an outcome than those treated with five days when the time to evaluation was 8 to 19 days, as opposed to 30 days (Pichichero 1997).

Potential biases in the review process

Potential weaknesses of meta-analysis techniques are that they incorporate existing biases and introduce new biases, some of which have predicted discordance of results between meta-analyses and single large RCTs (Borzak 1995; Egger 1997). To minimise bias during study selection, we used pre-determined inclusion criteria, and most trials were assessed in a blinded fashion, although recent evidence suggests that blinded evaluations are not necessary (Berlin 1997). We assessed publication bias in our funnel plot of sample size versus ORs (Higgins 2008) and with additional statistical tests (Duval 2000; Egger 1997) of significance. Publication bias was not evident. The issue of trial heterogeneity was addressed in our grouping of antibiotics according to pharmacokinetic profile (Schentag 1995).

In this review two studies (Gooch 1996; Hoberman 1997) contained two arms comparing longer course versus a short course of antibiotic. In the original review the short-course arm was counted twice against each long-course arm of that study. In the update of this review, we have modified our methods and combined both long-course arms in a single comparison against the short-course arm. In an effort to test whether this new method would change our results we compared the previous meta-analysis against a new analysis where we used the new statistical method but did not add the additional studies which are a part of this update. This new method did not have a great effect on our results.

Agreements and disagreements with other studies or reviews

Most of the 'short-acting' antibiotics in this study are commonly prescribed in primary practice and our sensitivity analyses indicated that comparisons between different short-acting antibiotics did not alter treatment outcomes. Comparability was also demonstrated between ceftriaxone and a longer course of antibiotics, al-

though the sample size of the consolidated trials was smaller. The equivalence observed between a three or five-day course of azithromycin and a 10-day course of other antibiotics was unchanged in sensitivity analyses, but these could not be performed for outcomes at 8 to 19 days. It is difficult to support the use of azithromycin or ceftriaxone based on their cost, and concern over indiscriminate use of broad-spectrum antibiotics (Rosenfeld 1996). Dagan et al documented increasing rates of erythromycin resistance to *Streptococcus pneumoniae* (*S. pneumoniae*) over the period that azithromycin was prescribed more often to children. The increased prescription of amoxicillin-clavulanate did alter resistance to penicillin (Dagan 2006). Ceftriaxone's intramuscular mode of administration may further limit its role (Eppesl 1997). Incidence of diarrhea and vomiting may be increased with antibiotic use in children with AOM (Del Mar 1997). Findings did not demonstrate that a shortened course of antibiotics decreased the likelihood of gastrointestinal effects, except when compared to a longer course of amoxicillin-clavulanate.

AUTHORS' CONCLUSIONS

Implications for practice

Evidence is increasing for a wait and watch approach to AOM. We believe that this is the most prudent approach for most children who are older than six months or do not have serious or complicated disease. If treatment is warranted, the clinician must decide if treatment for 7 to 10 days is worth the slightly reduced risk of treatment failure in the short term (< 21 days). Shorter courses can also be safely used, resulting in few side effects and, perhaps, a

lower risk of antibiotic resistant bacteria. Shorter courses may also be associated with higher levels of compliance.

Implications for research

In light of the increasing use of the wait and watch approach, it would be helpful to have trials that randomise participants to short versus long-course treatment with oral antibiotics after an unsuccessful observation period.

ACKNOWLEDGEMENTS

Support for this work was provided by the Canadian Cochrane Network and by the Department of Community Health Sciences at the University of Manitoba. We would like to acknowledge the previous review authors Hildes-Ripstein GE, Longstaffe SE, Wincott JL and Sitar DS for their work on the original document. Thanks to Mary Cheang, MSc for statistical assistance. In the update of this review thank you to Maria Ospina (MO) and Ben Vandermeer (BV) from the Alberta Research Centre for Health Evidence (ARCHE) for assistance with the risk of bias evaluation and statistical analysis. Also from ARCHE we would like to thank Nicola Hooton (NH) and Liza Bialy (LB) for assisting with screening, data extraction and author support; and librarians Carol Fiesen, Tamara Durec and Lisa Tjosvold. We would also like to thank Sarah Thorning, Acute Respiratory Tract Infections Group's Trials Search Co-ordinator who assisted with the last update and implemented the search strategy. Finally, we wish to thank the following people for commenting on the draft of this updated review: Dilip Raghavan, Sharon Sanders, Brian Westenberg, Max Bulsara and Peter Morris.

REFERENCES

References to studies included in this review

Adam 1996 {published data only}

Adam D. Five-day therapy with cefpodoxime vs. ten-day treatment with cefaclor in infants with acute otitis media. *Infection* 1995;**23**:398–9.

Adam 2000 {published data only}

Adam D. Efficacy and tolerability of 5-day vs. 10-day treatment with cefixime suspension in children with acute otitis media. *Drugs of Today* 2000;**36** (Suppl E):29–33.

Al-Ghamdi 1999 {published data only}

Al-Ghamdi YS, El-Kafrawi M, Aglan YI, Saleh MA. Efficacy of single-dose intramuscular ceftriaxone for treatment of acute otitis media in children. *Saudi Medical Journal* 1999; **20**(1):41–5.

Arguedas 1996 {published data only}

Arguedas A, Loaiza C, Herrera M, Mohs E. Comparative trial of 3-day azithromycin vs. 10-day amoxycillin/clavulante potassium in the treatment of children with acute otitis media with effusion. *International Journal of Antimicrobial Agents* 1996;**6**:233–8.

Arguedas 1997 {published data only}

Arguedas A, Loaiza C, Rodriguez F, Herrera ML, Mohs E. Comparative trial of 3 days of azithromycin versus 10 days of clarithromycin in the treatment of children with acute otitis media with effusion. *Journal of Chemotherapy* 1997;**9**: 44–50.

Arguedas 2005 {published data only}

Arguedas A, Emparanza P, Schwartz RH, Soley C, Guevara S, de Caprariis PJ, et al. A randomized, multicenter, double blind, double dummy trial of single dose azithromycin versus high dose amoxicillin for treatment of uncomplicated

- acute otitis media. *Pediatric Infectious Disease Journal* 2005; **24**(2):153–61.
- Aronovitz 1996** {published data only}
Aronovitz G. A multicenter, open label trial of azithromycin vs. amoxicillin/clavulanate for the management of acute otitis media in children. *Pediatric Infectious Disease Journal* 1996; **Suppl 15**:5–9.
- Arrieta 2003** {published data only}
Arrieta A, Arguedas A, Fernandez P, Block SL, Emperanza P, Vargas SL, et al. High-dose azithromycin versus high-dose amoxicillin-clavulanate for treatment of children with recurrent or persistent acute otitis media. *Antimicrobial Agents & Chemotherapy* 2003; **47**(10):3179–86.
- Barnett 1997** {published data only}
Barnett ED, Teele DW, Klein JO, Cabral HJ, Kharasch SJ. Comparison of ceftriaxone and trimethoprim-sulfamethoxazole for acute otitis media. *Pediatrics* 1997; **99**:23–8.
- Block 2000** {published data only}
Block SL, Kratzer J, Nemeth MA, Tack KJ. Five-day cefdinir course vs. ten-day cefprozil course for treatment of acute otitis media. *Pediatric Infectious Disease Journal* 2000; **19** (Suppl 12):147–52.
- Block 2003** {published data only}
Block SL, Arrieta A, Seibel M, McLinn S, Eppes S, Murphy MJ. Single-dose (30 mg/kg) azithromycin compared with 10-day amoxicillin/clavulanate (45 mg/kg per day) for the treatment of uncomplicated acute otitis media. *Current Therapeutic Research, Clinical & Experimental* 2003; **64** (Suppl A):30–42.
- Block 2004** {published data only}
Block SL, Busman TA, Paris MM, Bukofzer S. Comparison of five-day cefdinir treatment with ten-day low dose amoxicillin/clavulanate treatment for acute otitis media. *Pediatric Infectious Disease Journal* 2004; **23**(9):834–8.
- Boulesteix 1995** {published data only}
Boulesteix J, Dubreuil C, Moutot M, Rezvani Y, Rosembaum. Cefpodoxime proxetil five days versus cefixime eight days in the treatment of acute otitis media in children [Cefpodoxime proxetil 5 jours versus cefixime 8 jours, dans le traitement des otites moyennes aiguës de l'enfant]. *Médecine et Maladies Infectieuses* 1995; **25**:534–9.
- Catania 2004** {published data only}
Catania S, Gallo A. Clinical efficacy and tolerability of short course therapy with cefaclor compared with long-term therapy for treatment of acute otitis media in children [Il cefaclor nella terapia dell'otite media acuta in età pediatrica: confronto dell'efficacia e della tollerabilità di un trattamento breve rispetto a quello convenzionale]. *Le Infezioni in Medicina* 2004; **4**:259–65.
- Chamberlain 1994** {published data only}
Chamberlain JM, Boenning DA, Waisman Y, Ochenschlager DW, Klein BL. Single-dose ceftriaxone vs. 10 days of cefaclor for otitis media. *Clinical Pediatrics* 1994; **33**:642–6.
- Cohen 1997** {published data only}
Cohen R, de La Rocque F, Boucherat M, Levy C, Langue J, Bourrillon A. Randomized trial comparing 5-day cefpodoxime proxetil and 8-day amoxicillin-clavulanate treatment of acute otitis media in children [Etude randomisée cefpodoxime proxetil 5 jours versus amoxicilline-acide clavulanique 8 jours dans le traitement de l'otite moyenne aiguë de l'enfant]. *Médecine et Maladies Infectieuses* 1997; **27**:596–602.
- Cohen 1998** {published data only}
Cohen R, Levy C, Boucherat M, Langue J, de La Rocque F. A multicenter, randomized, double-blind trial of 5 versus 10 days of antibiotic therapy for acute otitis media in young children. *Journal of Pediatrics* 1998; **133**(5):634–9.
- Cohen 1999** {published data only}
Cohen R, Navel M, Grunberg J, Boucherat M, Geslin P, Derriennic M, et al. One dose ceftriaxone vs. ten days of amoxicillin/clavulanate therapy for acute otitis media: Clinical efficacy and change in nasopharyngeal flora. *Pediatric Infectious Disease Journal* 1999; **18**(5):403–9.
- Cohen 2000** {published data only}
Cohen R, Levy C, Boucherat M, Langue J, Autret E, Gehanno P, et al. Five vs. ten days of antibiotic therapy for acute otitis media in young children. *Pediatric Infectious Disease Journal* 2000; **19**(5):458–63.
- Dagan 2000a** {published data only}
Dagan R, Johnson CE, McLinn S, Abughali N, Feris J, Beibovitz E, et al. Bacteriologic and clinical efficacy of amoxicillin/clavulanate vs. azithromycin in acute otitis media. *Pediatric Infectious Disease Journal* 2000; **19**(2):95–104.
- Dagan 2000b** {published data only}
Dagan R, Leibovitz E, Fliss DM, Leiberman A, Jacobs MR, Craig W, et al. Bacteriologic efficacies of oral azithromycin and oral cefaclor in treatment of acute otitis media in infants and young children. *Antimicrobial Agents & Chemotherapy* 2000; **44**(1):43–50.
- Daniel 1993** {published data only}
Daniel RR. Comparison of azithromycin and co-amoxiclav in the treatment of otitis media in children. *Journal of Antimicrobial Chemotherapy* 1993; **31** (Suppl E):65–71.
- de Jose 1998** {published data only}
de Jose MI, Garcia de Miguel MJ, Aristegui J, Rodrigo C, del Castillo F, Borque C, et al. International multicenter study comparing azithromycin and amoxicillin-clavulanic acid in the treatment of pediatric acute otitis media [Estudio sobre la eficacia, seguridad y tolerancia de azitromicina frente a amoxicilina-clavulanico en el tratamiento de otitis media aguda]. *Acta Pediatrica Espanola* 1998; **56**(6):350–5.
- de Saintongue 1982** {published data only}
Chaput de Saintongue DM, Levine DF, Temple Savage I, Burgess GWS, Sharp J, Mayhew SR, et al. Trial of three-day and ten-day courses of amoxycillin in otitis media. *British Medical Journal* 1982; **284**:1078–81.

- Dunne 2003** {published data only}
Dunne MW, Latiolais T, Lewis B, Pistorius B, Bottenfield G, Moore WH, et al. Randomized, double-blind study of the clinical efficacy of 3 days of azithromycin compared with co-amoxiclav for the treatment of acute otitis media. *Journal of Antimicrobial Chemotherapy* 2003;**52**(3):469–72.
- Gooch 1996** {published data only}
Gooch WM, Blair E, Puopolo A, Paster ZR, Schwartz RH, Miller HC, et al. Effectiveness of five days of therapy with cefuroxime axetil suspension for treatment of acute otitis media. *Pediatric Infectious Disease Journal* 1996;**15**:157–64.
- Green 1993** {published data only}
Green SM, Rothrock SG. Single-dose intramuscular ceftriaxone for acute otitis media in children. *Pediatrics* 1993;**91**:23–30.
- Guven 2006** {published data only}
Guven M, Bulut Y, Sezer T, Aladag I, Eyibilen A, Etikan I. Bacterial etiology of acute otitis media and clinical efficacy of amoxicillin-clavulanate vs. azithromycin. *International Journal of Pediatric Otorhinolaryngology* 2006;**70**(5):915–23.
- Hendrickse 1988** {published data only}
Hendrickse WA, Kusmiesz H, Shelton S, Nelson JD. Five vs. ten days of therapy for acute otitis media. *Pediatric Infectious Disease Journal* 1988;**7**:14–23.
- Hoberman 1997** {published data only}
Hoberman A, Paradise JL, Burch DJ, Valinski WA, Hedrick JA, Aronovits GH, et al. Equivalent efficacy and reduced occurrence of diarrhea from a new formulation of amoxicillin/clavulanate potassium (Augmentin) for treatment of acute otitis media in children. *Pediatric Infectious Disease Journal* 1997;**16**(5):463–70.
- Hoberman 2005** {published data only}
Hoberman A, Dagan R, Leibovitz E, Rosenblut A, Johnson CE, Huff A, et al. Large dosage amoxicillin/clavulanate, compared with azithromycin, for the treatment of bacterial acute otitis media in children. *Pediatric Infectious Disease Journal* 2005;**24**(6):525–32.
- Ingvarsson 1982** {published data only}
Ingvarsson L, Lundgren K. Penicillin treatment of acute otitis media in children. *Acta Oto-Laryngologica* 1982;**94**:283–7.
- Kafetzis 1997** {published data only}
Kafetzis DA, Astra H, Mitropoulos L. Five-day versus ten-day treatment of acute otitis media with cefprozil. *European Journal of Clinical Microbiology & Infectious Diseases* 1997;**16**:283–6.
- Kara 1998** {published data only}
Kara CO, Ozuer MZ, Kilic I, Yalcin AN, Ergin H. Comparison of amoxicillin with second and third generation cephalosporins in the treatment of acute otitis media. *Infezioni in Medicina* 1998;**6**(2):93–5.
- Khurana 1996** {published data only}
Khurana CM. A multicenter, randomized, open label comparison of azithromycin and amoxicillin/clavulanate in acute otitis media among children attending day care or school. *Pediatric Infectious Disease Journal* 1996;**15**(Suppl 9):24–9.
- McLinn 1996** {published data only}
McLinn S. A multicenter, double blind comparison of azithromycin and amoxicillin/clavulanate for the treatment of acute otitis media in children. *Pediatric Infectious Disease Journal* 1996;**15**(Suppl 9):20–3.
- Meistrup-Larsen 1983** {published data only}
Meistrup-Larsen KI, Sorensen H, Johnsen NJ, Thomsen J, Mygind N, Sederberg-Olsen J. Two vs. seven days penicillin treatment for acute otitis media. *Acta Oto-Laryngologica* 1983;**96**:99–104.
- Mohs 1993** {published data only}
Mohs E, Rodriguez-Solares A, Rivas E, Hoshy ZE. A comparative study of azithromycin and amoxycillin in paediatric patients with acute otitis media. *Journal of Antimicrobial Chemotherapy* 1993;**31**(Suppl E):73–9.
- Oguz 2003** {published data only}
Oguz F, Unuvar E, Suoglu Y, Erdamar B, Dundar G, Katircioglu S, et al. Etiology of acute otitis media in childhood and evaluation of two different protocols of antibiotic therapy: 10 days cefaclor vs. 3 days azithromycin. *International Journal of Pediatric Otorhinolaryngology* 2003;**67**(1):43–51.
- Pessey 1999** {published data only}
Pessey JJ, Gehanno P, Thoroddsen E, Dagan R, Leibovitz E, Machac J, et al. Short course therapy with cefuroxime axetil for acute otitis media: results of a randomized multicenter comparison with amoxicillin/clavulanate. *Pediatric Infectious Disease Journal* 1999;**18**(10):854–9.
- Petalozza 1992** {published data only}
Petalozza G, Cioce C, Facchini M. Azithromycin in upper respiratory tract infections: a clinical trial in children with otitis media. *Scandinavian Journal of Infectious Diseases* 1992;**83**:22–5.
- Ploussard 1984** {published data only}
Ploussard JH. Evaluation of five days of cefaclor vs ten days of amoxicillin therapy in acute otitis media. *Current Therapeutic Research* 1984;**36**:641–5.
- Principi 1995** {published data only}
Principi N. Multicentre comparative study of the efficacy and safety of azithromycin compared with amoxicillin/clavulanic acid in the treatment of paediatric patients with otitis media. *European Journal of Clinical Microbiology & Infectious Diseases* 1995;**14**:669–7.
- Puczynski 1987** {published data only}
Puczynski MS, Stankiewicz JA, O'Keefe JP. Single dose amoxicillin treatment of acute otitis media. *Laryngoscope* 1987;**97**:16–8.
- Rodriguez 1996** {published data only}
Rodriguez AF. An open study to compare azithromycin with cefaclor in the treatment of children with acute otitis media. *Journal of Antimicrobial Chemotherapy* 1996;**37**(Suppl C):63–9.

Schaad 1993 {published data only}

Schaad UB. Multicentre evaluation of azithromycin in comparison with co-amoxiclav for the treatment of acute otitis media in children. *Journal of Antimicrobial Chemotherapy* 1993;**31**(Suppl E):81–8.

Varsano 1988 {published data only}

Varsano I, Frydman M, Amir J, Alpert G. Single intramuscular dose of ceftriaxone as compared to 7-day amoxicillin therapy for acute otitis media in children. *Chemotherapy* 1988;**34**(Suppl 1):39–46.

Varsano 1997 {published data only}

Varsano I, Volovitz B, Horev Z, Robinson J, Laks Y, Rosenbaum I, et al. Intramuscular ceftriaxone compared with oral amoxicillin-clavulanate for treatment of acute otitis media in children. *European Journal of Pediatrics* 1997;**156**(11):858–63.

Wang 2004 {published data only}

Wang CY, Lu CY, Hsieh YC, Lee CY, Huang LM. Intramuscular ceftriaxone in comparison with oral amoxicillin-clavulanate for the treatment of acute otitis media in infants and children. *Journal of Microbiology, Immunology & Infection* 2004;**37**(1):57–62.

References to studies excluded from this review

Bain 1985 {published data only}

Bain J, Murphy E, Ross F. Acute otitis media: clinical course among children who received a short course of high dose antibiotic. *British Medical Journal (Clinical Research Edition)* 1985;**291**:1243–6.

Gehanno 1990 {published data only}

Gehanno P, Taillebe M, Denis P, Jacquet P, Hoareau J, Gojon D, et al. Short-course cefotaxime compared with five-day co-amoxiclav in acute otitis media in children. *Journal of Antimicrobial Chemotherapy* 1990;**26**(Suppl A):29–36.

Harrison 1993 {published data only}

Harrison CJ, Chartrand SA, Pichichero ME. Microbiologic and clinical aspects of a trial of once daily cefixime compared with twice daily cefaclor for treatment of acute otitis media in infants and children. *Pediatric Infectious Disease Journal* 1993;**12**:62–9.

Jacobson 1979 {published data only}

Jacobson JA, Metcalf TJ, Parkin JL, Wenerstrom LG, Matsen JM. Evaluation of cefaclor and amoxycillin in the treatment of acute otitis media. *Postgraduate Medical Journal* 1979;**55**(Suppl 4):39–41.

Jenner 1987 {published data only}

Jenner PN, Tierney HE, Carle WK. A general practice study of a new approach to the use of antibiotics in the treatment of acute otitis media in children aged three and over. *British Journal of Clinical Practice* 1987;**41**:820–6.

Johnson 1991 {published data only}

Johnson CE, Carlin SA, Super DM, Rehms JM, Roberts DG, Christopher NC, et al. Cefixime compared with amoxicillin for treatment of acute otitis media. *Journal of Pediatrics* 1991;**119**:117–22.

Jones 1986 {published data only}

Jones R, Bain J. Three-day and seven-day treatment in acute otitis media: a double-blind antibiotic trial. *Journal of the Royal College of General Practitioners* 1986;**36**:356–8.

MacLoughlin 1996 {published data only}

MacLoughlin GJF, Barreto DG, de la Torre C, Pinetta EA, del Castillo F, Palma L. Cefpodoxime proxetil suspension compared with cefaclor suspension for treatment of acute otitis media in paediatric patients. *Journal of Antimicrobial Chemotherapy* 1996;**37**:565–73.

Mandel 1995 {published data only}

Mandel EM, Casselbrant ML, Rockette HE, Bluestone CD, Kurs-Lasky M. Efficacy of 20- vs. 10-day antimicrobial treatment for acute otitis media. *Pediatrics* 1995;**96**:5–13.

Murph 1993 {published data only}

Murph JR, Dusdieker LB, Booth B, Murph WE. Is treatment of acute otitis media with once-a-day amoxicillin feasible?. *International Pediatrics* 1993;**32**(9):528–34.

O'Doherty 1996 {published data only}

O'Doherty B. An open comparative study of azithromycin versus cefaclor in the treatment of patients with upper respiratory tract infections. *Journal of Antimicrobial Chemotherapy* 1996;**37**(Suppl C):71–81.

Roos 2000 {published data only}

Roos K, Larsson P. Efficacy of cefibuten in 5 vs 10 days treatment of recurrent acute otitis media in children. *International Journal of Pediatric Otorhinolaryngology* 2000;**55**:109–15.

Rubenstein 1965 {published data only}

Rubenstein MM, McBean JB, Hedgecock LD, Stickler GB. The treatment of acute otitis media in children. 3. A third clinical trial. *American Journal of Diseases of Children* 1965;**109**:308–13.

Scott 1990 {published data only}

Scott HV, Pannowitz D, Ketelbey JW. Cefixime: clinical trial against otitis media and tonsillitis. *New Zealand Medical Journal* 1990;**103**:25–6.

Simon 1997 {published data only}

Simon MW. Five- vs. 10-day treatment of acute otitis media with cefibuten in infants and children. *Advances in Therapy* 1997;**14**(6):312–7.

Spencer 1993 {published data only}

Spencer RC, Hannington J, Fraser S, Mason T. Cefpodoxime proxetil versus co-amoxiclav in the treatment of acute infections of the ear, nose and throat in children—a multicentre randomized study. Antibiotic Abstract 851. 18th International Congress of Chemotherapy (June 27 - July 2). Stockholm, Sweden, 1993:264.

Stickler 1964 {published data only}

Stickler GB, McBean JB. The treatment of acute otitis media in children. II. A second clinical trial. *JAMA* 1964;**187**:85–9.

Stickler 1967 {published data only}

Stickler GB, Rubenstein MM, McBean JB, Hedgecock LD, Hugstad A, Griffing T. Treatment of acute otitis media in

- children. IV. A fourth clinical trial. *American Journal of Diseases of Children* 1967;**114**:123–30.
- Suzuki 2009** {published data only}
Suzuki K, Baba S, Totsuka K, Hori S, Ubukata K, Nakashima M, et al. Double-blind comparative study of tebipenem pivoxil and high-dose cefditoren pivoxil in children with acute otitis media. *Japanese Journal of Chemotherapy* 2009;**57**(Suppl 1):167–85.
- Varsano 1994** {published and unpublished data}
Varsano I, Volovitz B, Horev Z, Robinson J, Laks Y, Rosenbaum I, et al. Single IM dose of ceftriaxone (CRO) compared to 10 days amoxicillin-clavulanate augmentin (AUG) for therapy of acute otitis media (AOM) in children. Sixth International Congress for Infectious Disease. Czech Republic, 1994; Vol. Session 117:1059.
- Additional references**
- Appelman 1991**
Appelman CLM, Claessen QPL, Touw-Otten FWMM, Hordijk GJ, de Melker RE. Co-amoxiclav in recurrent acute otitis media; placebo controlled study. *BMJ* 1991;**303**:1450–2.
- Berlin 1997**
Berlin JA. Does blinding of readers affect the results of meta-analyses. *Lancet* 1997;**350**:185–6.
- Berman 1995**
Berman S. Otitis media in developing countries. *Pediatrics* 1995;**96**:126–31.
- Boluyt 2008**
Boluyt N, Tjosvold L, Lefebvre C, Klassen T, Offringa M. Usefulness of systematic review search strategies in finding child health systematic reviews in MEDLINE. *Archives of Pediatrics and Adolescent Medicine* 2008;**162**(2):111–16.
- Borzak 1995**
Borzak S, Ridker PM. Discordance between meta-analyses and large-scale randomized, controlled trials. *Annals of Internal Medicine* 1995;**123**:873–7.
- Burke 1991**
Burke P, Bain J, Robinson D, Dunleavy J. Acute red ear in children: controlled trial of non-antibiotic treatment in general practice. *BMJ* 1991;**303**:558–62.
- Canafax 1991**
Canafax DM, Giebink GS. Clinical and pharmacokinetic basis for the antimicrobial treatment of acute otitis media. *Otolaryngologic Clinics of North America* 1991;**24**:859–74.
- Cohen 1992**
Cohen ML. Epidemiology of drug resistance: implications for a post-antimicrobial era. *Science* 1992;**257**:1050–5.
- Dagan 2006**
Dagan R, Barkai G, Leibovitz E, Dreifuss E, Greenberg D. Will reduction of antibiotic use reduce antibiotic resistance? The pneumococcus paradigm. *Pediatric Infectious Disease Journal* 2006;**25**(10):981–6.
- Del Mar 1997**
Del Mar C, Glasziou P, Hayem M. Are antibiotics indicated as initial treatment for children with acute otitis media? A meta-analysis. *BMJ* 1997;**314**:1526–9.
- DerSimonian 1986**
DerSimonian R, Laird N. Meta-analysis in clinical trials. *Controlled Clinical Trials* 1986;**7**:177–88.
- Duval 2000**
Duval S, Tweedie R. A nonparametric trim and fill method of accounting for publication bias in meta-analysis. *JAMA* 2000;**283**(449):89–98.
- Egger 1997**
Egger M, Smith GD, Schneider M. Bias in meta-analysis detected by a simple, graphical test. *BMJ* 1997;**315**:629–34.
- Eppes 1997**
Eppes SC, Klein JD, Lewis LL. Ceftriaxone for acute otitis media. *Pediatrics* 1997;**100**:157.
- Forgie 2009**
Forgie S, Zhanel G, Robinson J, Canadian Paediatric Society, Infectious Disease and Immunization Committee. Management of acute otitis media. Paediatric Child Health (Available at <http://www.cps.ca/ENGLISH/statements/ID/ID09-01.htm>) 2009; Vol. 14, issue 7:457–60.
- Froom 1990**
Froom J, Culpepper L, Grob P, Bartelds A, Bowers P, Bridges-Webb C, et al. Diagnosis and antibiotic treatment of acute otitis media: report from International Primary Care Network. *BMJ* 1990;**300**(6724):582–6.
- Gregoire 1995**
Gregoire G, Derderian F, Le Lorier J. Selecting the language of the publications included in a meta-analysis: is there a Tower of Babel bias?. *Journal of Clinical Epidemiology* 1995;**48**:159–63.
- Higgins 2008**
Higgins JPT, Green S (Editors). *Cochrane Handbook for Systematic Reviews of Interventions*. Chichester: John Wiley & Sons, 2008.
- Howie 1969**
Howie VM, Ploussard JH. The “in vitro sensitivity test” bacteriology of middle ear exudate. *Pediatrics* 1969;**44**:940–4.
- Kaleida 1991**
Kaleida PH, Casselbrant ML, Rockette HE, Paradise JL, Bluestone CD, Blatter MM, et al. Amoxicillin or myringotomy or both for acute otitis media: results of a randomized clinical trial. *Pediatrics* 1991;**87**:466–74.
- Koopman 2008**
Koopman L, Hoes AW, Glasziou PP, Appelman CL, Burke P, McCormick DP, et al. Antibiotic therapy to prevent the development of asymptomatic middle ear effusion in children with acute otitis media: a meta-analysis of individual patient data. *Archives of Otolaryngology - Head & Neck Surgery* 2008;**134**(2):128–32.

Kozyrskyj 1998

Kozyrskyj AL, Hildes-Ripstein GE, Longstaffe SEA, Wincott JL, Sitar DS, Klassen TP, et al. Treatment of acute otitis media with a shortened course of antibiotics. *JAMA* 1998;**279**(21):1736–42.

Laupacis 1988

Laupacis A, Sackett DL, Roberts RS. An assessment of clinically useful measures of the consequences of treatment. *New England Journal of Medicine* 1998;**318**:1728–33.

Lefebvre 2008

Lefebvre C, Eisinga A, McDonald S, Paul N. Enhancing access to reports of randomized trials published world-wide - the contribution of EMBASE records to the Cochrane Central Register of Controlled Trials (CENTRAL) in The Cochrane Library. *Emerging Themes in Epidemiology* 2008; **5**:13–25.

Murray 1994

Murray BE. Can antibiotic resistance be controlled?. *New England Journal of Medicine* 1994;**330**(17):1229–30.

Mygind 1981

Mygind N, Meistrup-Larsen KI, Thomsen J, Thomsem VF, Josefsson K, Sorensen H. Penicillin in acute otitis media: a double-blind placebo-controlled trial. *Clinical Otolaryngology* 1981;**6**:5–13.

Paradise 1995

Paradise JL. Managing otitis media: a time for change. *Pediatrics* 1995;**96**:712–5.

Paradise 1997

Paradise JL. Short-course antimicrobial treatment for acute otitis media. Not best for infants and young children. *JAMA* 1997;**278**:1640–2.

Pediatric Guidelines 2004

Subcommittee on Management of Acute Otitis Media, American Academy of Pediatrics, American Academy of Family Physicians. Clinical practice guidelines: diagnosis and management of acute otitis media. *Pediatrics* 2004;**113**(5):1451–65.

Pichichero 1997

Pichichero ME, Cohen R. Shortened course of antibiotic therapy for acute otitis media, sinusitis and tonsillopharyngitis. *Pediatric Infectious Disease Journal* 1997;**16**:680–95.

Rosenfeld 1994

Rosenfeld RM, Vertrees JE, Carr J, Cipolle RJ, Uden DL, Giebink GS, et al. Clinical efficacy of antimicrobial drugs for acute otitis media: meta-analysis of 5400 children from thirty-three randomized trials. *Journal of Pediatrics* 1994; **124**:355–67.

Rosenfeld 1996

Rosenfeld RM. An evidence-based approach to approach to treating otitis media. *Pediatric Clinics of North America* 1996;**43**:1165–81.

Rovers 2004

Rovers MM, Schilder AGM, Zielhuis GA, Rosenfeld RM. Otitis media. *The Lancet* 2004;**363**:465–73.

Sanders 2009

Sanders S, Glasziou PP, Del Mar C, Rovers M. Antibiotics for acute otitis media in children. *Cochrane Database of Systematic Reviews* 2009, Issue 2. [DOI: 10.1002/14651858.CD000219.pub2]

Schentag 1995

Schentag JJ. Antibiotic treatment of acute otitis media in children: dosing considerations. *Pediatric Infectious Disease Journal* 1995;**14**:S30–3.

Spiro 2008

Spiro DM, Arnold DH. The concept and practice of a wait-and-see approach to acute otitis media. *Current Opinion in Pediatrics* 2008;**20**:72–8.

Thompson 1994

Thompson SG. Why sources of heterogeneity in meta-analysis should be investigated. *BMJ* 1994;**309**:1351–5.

van Buchem 1981

van Buchem FL, Dunk JHM, Van't Hof MA. Therapy of acute otitis media: myringotomy, antibiotic or neither. *Lancet* 1981;**II**:883–7.

Vouloumanou 2009

Vouloumanou EK, Karageorgopoulos DE, Kazantzi MS, Kapaskelis AM, Falagas ME. Antibiotics versus placebo or watchful waiting for acute otitis media: a meta-analysis of randomized controlled trials. *Journal of Antimicrobial Chemotherapy* 2009;**64**:16–24.

References to other published versions of this review**Kozyrski 1998**

Kozyrski AL, Hildes-Ripstein E, Longstaffe SA, Wincott JL, Sitar DS, Klassen TP, et al. Treatment of acute otitis media with a shortened course of antibiotics. *JAMA* 1998;**279**(21):1736–42.

Kozyrskyj 2000

Kozyrskyj AL, Hildes-Ripstein GE, Longstaffe SEA, Wincott JL, Sitar DS, Klassen TP, et al. Short course antibiotics for acute otitis media. *Cochrane Database of Systematic Reviews* 2000, Issue 2. [DOI: 10.1002/14651858.CD001095]

* Indicates the major publication for the study

CHARACTERISTICS OF STUDIES

Characteristics of included studies [ordered by study ID]

Adam 1996

Methods	RCT	
Participants	German multicentre trial of children 3 months to 6 years old	
Interventions	Cefpodoxime 40 mg to 60 mg twice daily for 5 days versus cefaclor 40 mg/kg/day 3 times daily for 10 days	
Outcomes	Absence or improvement of signs and symptoms at 3 days and no relapse at 3 weeks after study entry	
Notes	-	
<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement
Blinding (performance bias and detection bias) Treatment failure at 1 month or less	High risk	
Incomplete outcome data (attrition bias) Treatment failure at 1 month or less	Low risk	
Selective reporting (reporting bias)	Low risk	
Other bias	Unclear risk	Funding not reported

Adam 2000

Methods	Double-blind RCT	
Participants	German children aged 2 to 14 years	
Interventions	Cefixime 8 mg/kg/day for 5 days versus same treatment for 10 days	
Outcomes	Complete recovery or improvement of symptoms by day 11, clinical response rate at day 6, rate of relapse at day 28	
Notes	-	

<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement
Blinding (performance bias and detection bias) Treatment failure at 1 month or less	Low risk	
Incomplete outcome data (attrition bias) Treatment failure at 1 month or less	High risk	
Selective reporting (reporting bias)	High risk	
Other bias	Unclear risk	Funding not reported

Al-Ghamdi 1999

Methods	RCT
Participants	Saudi children aged 6 months to 6 years
Interventions	Single intramuscular dose 50 mg/kg ceftriaxone versus 40 mg/kg/day Augmentin 3 times daily for 10 days
Outcomes	Resolution of symptoms at 10 days after study entry; improved otoscopic and tympanometric measurements at 60 days after study entry
Notes	-

<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement
Incomplete outcome data (attrition bias) Treatment failure at 1 month or less	Low risk	
Selective reporting (reporting bias)	Low risk	
Other bias	Unclear risk	Did not report baselines or funding

Arguedas 1996

Methods	RCT
Participants	Children 6 months to 12 years old in an ambulatory unit in Costa Rica
Interventions	Azithromycin 10 mg/kg once daily for 3 days versus amoxil-clavulanate 40 mg/kg/day 3 times daily for 10 days
Outcomes	Absence or improvement of signs and symptoms at 11 to 13 days, 28 to 30 days and 55 to 60 days after study entry
Notes	-

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	
Allocation concealment (selection bias)	Low risk	
Blinding (performance bias and detection bias) Treatment failure at 1 month or less	High risk	
Incomplete outcome data (attrition bias) Treatment failure at 1 month or less	Low risk	
Other bias	Low risk	Industry funding

Arguedas 1997

Methods	RCT
Participants	Children 6 months to 12 years old seen at an ambulatory unit in Costa Rica
Interventions	Azithromycin 10 mg/kg once daily for 3 days versus clarithromycin 15 mg/kg/day twice daily for 10 days
Outcomes	Resolution of symptoms with or without the presence of middle ear fluid at 10 to 11 days and 28 to 32 days after study entry
Notes	-

Risk of bias

Arguedas 1997 (Continued)

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	
Allocation concealment (selection bias)	Low risk	
Blinding (performance bias and detection bias) Treatment failure at 1 month or less	High risk	
Incomplete outcome data (attrition bias) Treatment failure at 1 month or less	Low risk	
Other bias	Low risk	Industry funding

Arguedas 2005

Methods	Double-blind, double-dummy multinational RCT
Participants	Children 6 to 30 months of age in multiple countries
Interventions	Azithromycin single dose 30 mg/kg versus amoxicillin 90 mg/kg/day for 10 days in 2 divided doses
Outcomes	Absence or improvement of symptoms at 12 to 14 days and at end of study (day 25 to 28)
Notes	-

Risk of bias

Bias	Authors' judgement	Support for judgement
Blinding (performance bias and detection bias) Treatment failure at 1 month or less	Low risk	
Incomplete outcome data (attrition bias) Treatment failure at 1 month or less	Low risk	
Selective reporting (reporting bias)	High risk	
Other bias	High risk	Interim analysis, industry funding

Aronovitz 1996

Methods	RCT
Participants	US multicentre trial of children 2 to 15 years old
Interventions	Azithromycin 5 to 10 mg/kg once daily for 5 days versus amoxil-clavulanate 40 mg/kg/day 3 times daily for 10 days
Outcomes	Absence or improvement of signs and symptoms at 11 days and 30 days after study entry
Notes	-

Risk of bias

Bias	Authors' judgement	Support for judgement
Blinding (performance bias and detection bias) Treatment failure at 1 month or less	High risk	
Incomplete outcome data (attrition bias) Treatment failure at 1 month or less	High risk	
Selective reporting (reporting bias)	Low risk	
Other bias	Unclear risk	No funding declared

Arrieta 2003

Methods	Double-blind, double-dummy, multicentre trial
Participants	US and Latin American children aged 6 months to 6 years
Interventions	Azithromycin 20 mg/kg/day once daily for 3 days versus amoxicillin/clavulanate 45/6.4 mg/kg/day twice daily for 10 days plus additional amoxil 45 mg/kg/day twice daily for 10 days
Outcomes	Clinical response (improvement or cure) among clinically evaluated patients at day 28 to 32; clinical response at day 12 to 16
Notes	-

Risk of bias

Bias	Authors' judgement	Support for judgement
------	--------------------	-----------------------

Arrieta 2003 (Continued)

Blinding (performance bias and detection bias) Treatment failure at 1 month or less	Low risk	
Incomplete outcome data (attrition bias) Treatment failure at 1 month or less	Low risk	
Selective reporting (reporting bias)	Low risk	
Other bias	Unclear risk	Corresponding author from Pfizer, unclear independence of authors, industry funding

Barnett 1997

Methods	RCT
Participants	Boston, US multicentre trial of children 3 months to 3 years old
Interventions	Ceftriaxone 50 mg/kg intramuscularly x one dose versus trimethoprim-sulfamethoxazole twice daily for 10 days
Outcomes	Absence of signs and symptoms at 14 days and 28 days after study entry
Notes	-

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	
Blinding (performance bias and detection bias) Treatment failure at 1 month or less	High risk	Personnel not blinded
Incomplete outcome data (attrition bias) Treatment failure at 1 month or less	High risk	
Selective reporting (reporting bias)	Low risk	
Other bias	Unclear risk	Industry funding

Block 2000

Methods	Comparative, randomized, investigator blinded multicentre trial
Participants	US children aged 6 months through 12 years
Interventions	Cefdinir 14 mg/kg/day twice daily for 5 days versus cefprozil 30 mg/kg twice daily for 10 days
Outcomes	Cure or failure at end of treatment visit, cure 11 to 16 days post-therapy, long-term follow-up days 38 to 45 after study entry
Notes	-

Risk of bias

Bias	Authors' judgement	Support for judgement
Blinding (performance bias and detection bias) Treatment failure at 1 month or less	High risk	Only investigators blinded
Other bias	Low risk	Industry funded

Block 2003

Methods	Double-blind, placebo-controlled, RCT
Participants	US children aged 6 months to 12 years
Interventions	Single dose of 30 mg/kg azithromycin versus amoxicillin/clavulanate 45 mg/kg twice daily for 10 days
Outcomes	Absence or improvement of signs and symptoms on day 12 to 16 and day 28 to 32
Notes	-

Risk of bias

Bias	Authors' judgement	Support for judgement
Blinding (performance bias and detection bias) Treatment failure at 1 month or less	Low risk	
Incomplete outcome data (attrition bias) Treatment failure at 1 month or less	Low risk	

Block 2003 (Continued)

Selective reporting (reporting bias)	Low risk	
Other bias	Unclear risk	Industry funding

Block 2004

Methods	Randomised, investigator blinded multicentre study
Participants	US children aged 6 months through 6 years
Interventions	Cefdinir 14 mg/kg twice daily for 5 days versus amoxicillin/clavulanate 45/6.4 mg/kg twice daily for 10 days
Outcomes	Cure at end of treatment (7 to 9 days or 12 to 14 days) and days 25 to 28 of the study
Notes	-

Risk of bias

Bias	Authors' judgement	Support for judgement
Blinding (performance bias and detection bias) Treatment failure at 1 month or less	High risk	Only investigator blinded
Selective reporting (reporting bias)	High risk	
Other bias	Low risk	Industry funding

Boulesteix 1995

Methods	RCT
Participants	French multicentre trial of children 6 months to 6 years old
Interventions	Cefpodoxime 4 mg/kg twice daily for 5 days versus cefixime 4 mg/kg twice daily for 8 days
Outcomes	Absence or improvement of signs and symptoms at 8 to 10 days and no recurrence at 30 to 40 days after study entry
Notes	-

Risk of bias

Boulesteix 1995 (Continued)

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	
Blinding (performance bias and detection bias) Treatment failure at 1 month or less	High risk	
Incomplete outcome data (attrition bias) Treatment failure at 1 month or less	Low risk	
Selective reporting (reporting bias)	Low risk	
Other bias	Unclear risk	Funding not declared

Catania 2004

Methods	RCT
Participants	Italian children 2 to 6 years
Interventions	Cefaclor 40 mg/kg/day for 5 days versus cefaclor 40 mg/kg/day for 10 days
Outcomes	Clinical improvement or failure at end of therapy and 15 to 20 days after end of therapy. Relapse at 15 to 20 days after end of therapy
Notes	Italian

Risk of bias

Bias	Authors' judgement	Support for judgement
Incomplete outcome data (attrition bias) Treatment failure at 1 month or less	Low risk	
Selective reporting (reporting bias)	Low risk	
Other bias	Unclear risk	No funding declared

Chamberlain 1994

Methods	RCT
Participants	Children 18 months to 6 years old in Washington, DC

Chamberlain 1994 (Continued)

Interventions	Ceftriaxone 50 mg/kg intramuscularly x one dose versus cefaclor 40 mg/kg/day for 10 days
Outcomes	Absence or improvement of symptoms, and normal or improved tympanogram at 7 to 10 days and monthly for 90 days after study entry
Notes	-

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	
Blinding (performance bias and detection bias) Treatment failure at 1 month or less	Low risk	
Selective reporting (reporting bias)	Low risk	
Other bias	Low risk	Industry funding

Cohen 1997

Methods	RCT
Participants	French multicentre trial of children 4 months to 3 years old
Interventions	Cefpodoxime 8 mg/kg/day twice daily for 5 days versus amoxil-clavulanate 80 mg/kg/day 3 times daily for 8 days
Outcomes	Absence or improvement of signs and symptoms and no additional antibiotics at 9 to 14 days and 20 to 30 days after study entry
Notes	-

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	
Blinding (performance bias and detection bias) Treatment failure at 1 month or less	High risk	

Cohen 1997 (Continued)

Incomplete outcome data (attrition bias) Treatment failure at 1 month or less	Low risk	
Selective reporting (reporting bias)	Low risk	
Other bias	Unclear risk	Funding not declared

Cohen 1998

Methods	Randomised controlled, double-blind trial
Participants	French trial of children aged 4 to 30 months
Interventions	Amoxicillin/clavulanate 80/10 mg/kg/day 3 times daily for 5 days versus same treatment for 10 days
Outcomes	Resolution or improvement of signs and symptoms at 12 to 14 days and 28 to 42 days after study entry
Notes	-

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	
Blinding (performance bias and detection bias) Treatment failure at 1 month or less	Low risk	
Selective reporting (reporting bias)	High risk	
Other bias	Unclear risk	The only significant difference at baseline was diarrhea, but GI symptoms are an outcome; industry funding

Cohen 1999

Methods	Randomised open trial
Participants	French trial of children aged 4 to 30 months

Cohen 1999 (Continued)

Interventions	Ceftriaxone 50 mg/kg intramuscularly x one dose versus amoxicillin/clavulanate 80/10 mg/kg/day 3 times daily for 10 days
Outcomes	Resolution or improvement of signs and symptoms at 12 to 14 days and 28 to 42 days after study entry
Notes	-

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	
Blinding (performance bias and detection bias) Treatment failure at 1 month or less	High risk	
Incomplete outcome data (attrition bias) Treatment failure at 1 month or less	Low risk	Unclear for the 28 to 42-day follow-up period
Selective reporting (reporting bias)	Low risk	
Other bias	Low risk	Industry funding

Cohen 2000

Methods	Randomised controlled, double-blind trial
Participants	French trial of children aged 4 to 30 months
Interventions	Cefpodoxime proxetil 8 mg/kg/day for 5 days versus same regimen for 10 days
Outcomes	Resolution or improvement of signs and symptoms at 12 to 14 days and 28 to 42 days after study entry
Notes	-

Risk of bias

Bias	Authors' judgement	Support for judgement
------	--------------------	-----------------------

Cohen 2000 (Continued)

Random sequence generation (selection bias)	Low risk	
Blinding (performance bias and detection bias) Treatment failure at 1 month or less	Low risk	
Selective reporting (reporting bias)	Low risk	
Other bias	Low risk	Industry funding

Dagan 2000a

Methods	Single-blind, RCT
Participants	Multicentre trial (Israel, US, Dominican Republic) of children aged 6 to 48 months
Interventions	Azithromycin, 10 mg/kg on day 1 then 5 mg/kg on days 2 to 5 versus amoxicillin/clavulanate 45/6.4 mg/kg in 2 divided doses for 10 days
Outcomes	Resolution of signs and symptoms on day 12 to 14 and day 22 to 28, bacteriologic cure days 4 to 6
Notes	

Risk of bias

Bias	Authors' judgement	Support for judgement
Incomplete outcome data (attrition bias) Treatment failure at 1 month or less	High risk	
Selective reporting (reporting bias)	Low risk	
Other bias	Low risk	Industry funding

Dagan 2000b

Methods	Open label, RCT
Participants	Israeli trial of children aged 3 to 36 months old
Interventions	Azithromycin 10 mg/kg/day for 3 days versus cefaclor 40 mg/kg/day in 3 divided doses for 10 days

Dagan 2000b (Continued)

Outcomes	Resolution or improvement of signs and symptoms on day 10 and day 17 +/- 2	
Notes	-	
<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement
Blinding (performance bias and detection bias) Treatment failure at 1 month or less	Low risk	
Selective reporting (reporting bias)	Low risk	
Other bias	Low risk	Industry funding

Daniel 1993

Methods	RCT	
Participants	European multicentre trial of children 2 to 8 years old	
Interventions	Azithromycin 10 mg/kg daily for 3 days versus amoxil-clavulanate 3 times daily for 10 days	
Outcomes	Absence or improvement of signs and symptoms at 10 to 12 days after study entry	
Notes	-	
<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement
Blinding (performance bias and detection bias) Treatment failure at 1 month or less	High risk	
Incomplete outcome data (attrition bias) Treatment failure at 1 month or less	Low risk	
Selective reporting (reporting bias)	Low risk	
Other bias	High risk	Potential bias in that in the short course, azithromycin patients were placed on different antibiotics if failing, whereas for co-amoxil they stayed on co-amoxil; industry funding

de Jose 1998

Methods	RCT
Participants	Spanish children 6 months to 12 years old
Interventions	Azithromycin 10 mg/kg/day once daily for 3 days versus amoxicillin/clavulanate 40 mg/kg/day 3 times daily for 10 days
Outcomes	Clinical cure, improvement or failure at end of treatment
Notes	Spanish

Risk of bias

Bias	Authors' judgement	Support for judgement
Blinding (performance bias and detection bias) Treatment failure at 1 month or less	High risk	Outcome assessors not blinded
Incomplete outcome data (attrition bias) Treatment failure at 1 month or less	Low risk	
Selective reporting (reporting bias)	Low risk	
Other bias	Unclear risk	Funding not reported

de Saintongue 1982

Methods	Randomised controlled, double-blind trial
Participants	British multicentre trial of children 2 to 10 years old
Interventions	Amoxicillin 125/250 mg 3 times daily for 3 days + placebo for 7 days versus amoxicillin 125/250 mg 3 times daily for 10 days
Outcomes	Resolution of signs and symptoms at 13 to 16 days after and recurrence within 12 weeks of study entry
Notes	-

Risk of bias

Bias	Authors' judgement	Support for judgement
Blinding (performance bias and detection bias) Treatment failure at 1 month or less	Low risk	

de Saintongue 1982 (Continued)

Incomplete outcome data (attrition bias) Treatment failure at 1 month or less	Low risk	
Selective reporting (reporting bias)	Low risk	
Other bias	Low risk	

Dunne 2003

Methods	Randomised controlled double-blind trial
Participants	US trial of children 6 months to 12 years
Interventions	Azithromycin 10 mg/kg/day for 3 days versus co-amoxiclav 45 mg/kg/day twice a day for 10 days
Outcomes	Absence of symptoms at day 24 to 28, absence or improvement of symptoms at day 10
Notes	-

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	
Blinding (performance bias and detection bias) Treatment failure at 1 month or less	Unclear risk	Method not described
Incomplete outcome data (attrition bias) Treatment failure at 1 month or less	Low risk	
Selective reporting (reporting bias)	Low risk	
Other bias	Unclear risk	Industry funding, first author from Pfizer

Gooch 1996

Methods	Randomised controlled, double-blind trial
Participants	US multicentre trial of children 3 months to 12 years old
Interventions	Cefuroxime 30 mg/kg/day twice daily for 5 days + placebo twice daily for 5 days versus cefuroxime 30 mg/kg/day twice daily for 10 days

Gooch 1996 (Continued)

Outcomes	Absence or improvement of signs and symptoms at 14 to 18 days after study entry	
Notes	-	
<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement
Incomplete outcome data (attrition bias) Treatment failure at 1 month or less	High risk	
Selective reporting (reporting bias)	Low risk	
Other bias	Low risk	Industry funding

Green 1993

Green 1998

Methods	Randomised controlled, double-blind trial	
Participants	Children 5 months to 5 years old seen in emergency departments or medical centres in California, North Carolina, US	
Interventions	Ceftriaxone 50 mg/kg intramuscularly x 1 dose + placebo 3 times daily for 10 days versus placebo intramuscularly x 1 dose + amoxicillin 40 mg/kg 3 times daily for 10 days	
Outcomes	Absence of symptoms at 10 days and no recurrence at 11 to 30 days and 31 to 90 days after study entry	
Notes	-	
<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	
Blinding (performance bias and detection bias) Treatment failure at 1 month or less	Low risk	
Selective reporting (reporting bias)	Low risk	
Other bias	Low risk	Industry funding

Güven 2006

Methods	RCT
Participants	Turkish children between 6 months and 12 years
Interventions	Azithromycin (10 mg/kg/24 hours, per oral) once daily for 3 days versus amoxicillin-clavulanate (45 mg/kg/24 hours to 6.4 mg/kg/24 hours, peroral in 2 divided doses) for 10 days
Outcomes	Clinical cure, failure or improvement at the end of treatment and clinical cure, failure, improvement, relapse or re-infection at 26 to 28 days following study entry
Notes	-

Risk of bias

Bias	Authors' judgement	Support for judgement
Blinding (performance bias and detection bias) Treatment failure at 1 month or less	Unclear risk	Regimen was not equivalent
Incomplete outcome data (attrition bias) Treatment failure at 1 month or less	Low risk	
Selective reporting (reporting bias)	Low risk	
Other bias	Low risk	Funding not declared

Hendrickse 1988

Methods	Randomised controlled, double-blind trial
Participants	Children 1 month to 12 years old seen at a medical centre in Dallas, US
Interventions	Cefaclor 40 mg/kg/day twice daily for 5 days + placebo for 5 days versus cefaclor 40 mg/kg/day twice daily for 10 days
Outcomes	Resolution of middle ear fluid, or healing of tympanic membrane with resolution of signs and symptoms at 10 days, 30 days, 60 days and 90 days after study entry
Notes	-

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	

Hendrickse 1988 (Continued)

Blinding (performance bias and detection bias) Treatment failure at 1 month or less	Low risk	
Incomplete outcome data (attrition bias) Treatment failure at 1 month or less	Low risk	
Selective reporting (reporting bias)	Low risk	
Other bias	Low risk	Industry funding

Hoberman 1997

Methods	RCT
Participants	US and Canadian multicentre trial of children 2 months to 12 years old
Interventions	Amoxil-clavulanate (new formulation) twice daily for 5 days versus amoxil-clavulanate (new formulation) twice daily for 10 days or amoxil-clavulanate (old formulation) 3 times daily for 10 days
Outcomes	Absence or improvement of signs and symptoms with or without middle ear effusion at 12 to 14 days and 32 to 38 days after study entry
Notes	-

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	
Blinding (performance bias and detection bias) Treatment failure at 1 month or less	Unclear risk	Only investigators blinded
Incomplete outcome data (attrition bias) Treatment failure at 1 month or less	Low risk	
Selective reporting (reporting bias)	Low risk	
Other bias	Unclear risk	Industry funding and authors from SmithKline Beecham, baseline differences in exposure to cigarette smoke

Hoberman 2005

Methods	Investigator blinded, randomized controlled multicentre trial
Participants	Multicentre trial of children 6 to 30 months of age in various countries
Interventions	Azithromycin 10 mg/kg for 1 day followed by 5 mg/kg/d for 5 days versus amoxicillin/clavulanate 90/6.4 mg/kg/d in 2 divided doses for 10 days
Outcomes	Absence or improvement of signs and symptoms at day 4 to 6, 12 to 14 and 21 to 25 days after study entry
Notes	-

Risk of bias

Bias	Authors' judgement	Support for judgement
Blinding (performance bias and detection bias) Treatment failure at 1 month or less	Unclear risk	Investigator blinded
Selective reporting (reporting bias)	Low risk	
Other bias	Low risk	Industry funding

Ingvarsson 1982

Methods	RCT
Participants	Swedish children 6 months to 7 years old
Interventions	Penicillin-V 25 mg/kg twice daily for 5 days versus penicillin-V 25 mg/kg twice daily for 10 days
Outcomes	Improved signs and symptoms at 10 to 12 days and normal eardrum at 28 to 30 days after study entry
Notes	-

Risk of bias

Bias	Authors' judgement	Support for judgement
Blinding (performance bias and detection bias) Treatment failure at 1 month or less	High risk	Not mentioned and no placebo

Ingvarsson 1982 (Continued)

Incomplete outcome data (attrition bias) Treatment failure at 1 month or less	Low risk	
Selective reporting (reporting bias)	Low risk	
Other bias	Unclear risk	Funding not reported

Kafetzis 1997

Methods	RCT
Participants	Greek multicentre trial of children 2 to 172 months old
Interventions	Cefprozil 30 mg/kg/day twice daily for 5 days versus cefprozil 30 mg/kg/day twice daily for 10 days
Outcomes	Absence or improvement of signs and symptoms at 7 to 14 days and 28 to 32 days after study entry
Notes	

Risk of bias

Bias	Authors' judgement	Support for judgement
Blinding (performance bias and detection bias) Treatment failure at 1 month or less	High risk	
Incomplete outcome data (attrition bias) Treatment failure at 1 month or less	Low risk	
Selective reporting (reporting bias)	Low risk	
Other bias	Unclear risk	Funding not reported

Kara 1998

Methods	RCT
Participants	Turkish children aged 6 months to 6 years
Interventions	Ceftriaxone 50 mg/kg single intramuscular dose versus cefuroxime axetil 30 mg/kg 3 times daily for 10 days versus

Kara 1998 (Continued)

	amoxicillin 40 mg/kg 3 times daily for 10 days	
Outcomes	Clinical cure rate at 10 days after study entry	
Notes	-	
<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement
Other bias	Unclear risk	Funding not reported, baseline characteristics not reported

Khurana 1996

Methods	RCT	
Participants	US multicentre trial of children 6 months to 12 years old	
Interventions	Azithromycin 5 to 10 mg/kg once daily for 5 days versus amoxil-clavulanate 40 mg/kg/day 3 times daily for 10 days	
Outcomes	Absence or improvement of signs and symptoms at 45 days after study entry	
Notes	-	
<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement
Blinding (performance bias and detection bias) Treatment failure at 1 month or less	High risk	
Incomplete outcome data (attrition bias) Treatment failure at 1 month or less	Low risk	
Selective reporting (reporting bias)	Low risk	
Other bias	Unclear risk	Funding not reported

McLinn 1996

Methods	Randomised controlled, double-blind trial
Participants	US multicentre trial of children 1 to 15 years old
Interventions	Azithromycin 5 to 10 mg/kg once daily/placebo twice daily for 5 days + placebo 3 times daily for 5 days versus amoxil-clavulanate 40 mg/kg/day 3 times daily for 10 days
Outcomes	Absence or improvement of signs and symptoms at 11 days and 30 days after study entry
Notes	-

Risk of bias

Bias	Authors' judgement	Support for judgement
Blinding (performance bias and detection bias) Treatment failure at 1 month or less	Low risk	
Incomplete outcome data (attrition bias) Treatment failure at 1 month or less	High risk	
Selective reporting (reporting bias)	Low risk	
Other bias	Unclear risk	Funding not reported

Meistrup-Larsen 1983

Methods	Randomised controlled, double-blind trial
Participants	Children 1 to 10 years old in Copenhagen
Interventions	Penicillin-V 55 mg/kg twice daily for 2 days + placebo for 5 days versus penicillin-V 55 mg/kg twice daily for 7 days
Outcomes	Absence of symptoms and no otorrhoea or contralateral otitis at 6 to 7 days after study entry
Notes	-

Risk of bias

Bias	Authors' judgement	Support for judgement
------	--------------------	-----------------------

Meistrup-Larsen 1983 (Continued)

Blinding (performance bias and detection bias) Treatment failure at 1 month or less	Low risk	
Incomplete outcome data (attrition bias) Treatment failure at 1 month or less	Low risk	
Selective reporting (reporting bias)	Low risk	
Other bias	Unclear risk	Funding not reported, doses prepared in the home

Mohs 1993

Methods	Comparative, open trial (randomisation not mentioned)
Participants	Multicentre trial (South America and Egypt) of children 2 to 12 years old
Interventions	Azithromycin 10 mg/kg once daily for 3 days versus amoxicillin 10 mg/kg 3 times daily for 10 days
Outcomes	Absence or improvement of signs and symptoms at 11 to 13 days after study entry
Notes	-

Risk of bias

Bias	Authors' judgement	Support for judgement
Blinding (performance bias and detection bias) Treatment failure at 1 month or less	High risk	
Incomplete outcome data (attrition bias) Treatment failure at 1 month or less	Low risk	
Selective reporting (reporting bias)	Low risk	
Other bias	Unclear risk	Funding not reported

Oguz 2003

Methods	Randomised, single-blind, open study
Participants	Turkish children aged 6 months to 12 years
Interventions	Azithromycin 10 mg/kg/day for 3 days versus cefaclor 40 mg/kg/day in 3 divided doses for 10 days
Outcomes	Resolution of all clinical and otoscopic findings at 3 to 5, 10 and 30 days after study entry; relapse at 10 and 30 days
Notes	-

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	
Blinding (performance bias and detection bias) Treatment failure at 1 month or less	Low risk	
Incomplete outcome data (attrition bias) Treatment failure at 1 month or less	Low risk	
Selective reporting (reporting bias)	Low risk	
Other bias	Unclear risk	Funding not reported

Pessey 1999

Methods	Randomised, open, multinational study
Participants	Children aged 6 to 36 months
Interventions	Cefuroxime axetil 30 mg/kg/day twice daily for 5 days versus amoxicillin/clavulanate 40 mg/kg/day 3 times daily for 10 days versus amoxicillin/clavulanate 80 mg/kg/day 3 times daily for 8 days
Outcomes	Absence of symptoms 1 to 4 days after therapy completion; no relapse 21 to 28 days after therapy completion
Notes	-

Risk of bias

Pessey 1999 (Continued)

Bias	Authors' judgement	Support for judgement
Blinding (performance bias and detection bias) Treatment failure at 1 month or less	High risk	Open study
Incomplete outcome data (attrition bias) Treatment failure at 1 month or less	Low risk	
Selective reporting (reporting bias)	Low risk	
Other bias	Unclear risk	Corresponding author from Glaxo, funding not reported

Petalozza 1992

Methods	RCT
Participants	Italian children 11 months to 9 years old
Interventions	Azithromycin 10 mg/kg once daily for 3 days versus amoxil-clavulanate 50 mg/kg/day twice daily for 10 days
Outcomes	Absence or improvement of signs and symptoms at 12 to 14 days and 30 days after study entry
Notes	-

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	High risk	Openly randomized
Blinding (performance bias and detection bias) Treatment failure at 1 month or less	High risk	
Incomplete outcome data (attrition bias) Treatment failure at 1 month or less	Low risk	
Other bias	Unclear risk	Funding not reported

Ploussard 1984

Methods	RCT
Participants	Children 5 months to 5 years old in Alabama, US
Interventions	Cefaclor 40 mg/kg 3 times daily for 5 days versus amoxicillin 40 mg/kg 3 times daily for 10 days
Outcomes	Absence or improvement of signs and symptoms at 10 to 16 days after study entry
Notes	-

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	
Blinding (performance bias and detection bias) Treatment failure at 1 month or less	High risk	
Incomplete outcome data (attrition bias) Treatment failure at 1 month or less	Low risk	
Selective reporting (reporting bias)	Low risk	
Other bias	Unclear risk	Funding not reported

Principi 1995

Methods	RCT
Participants	Multicentre trial (Europe, South America) of children 6 months to 12 years old
Interventions	Azithromycin 10 mg/kg once daily for 3 days versus amoxil-clavulanate 40 mg/kg 3 times daily for 10 days
Outcomes	Absence or improvement of signs and symptoms at 10 to 14 days after study entry
Notes	-

Risk of bias

Bias	Authors' judgement	Support for judgement
------	--------------------	-----------------------

Principi 1995 (Continued)

Blinding (performance bias and detection bias) Treatment failure at 1 month or less	High risk	
Incomplete outcome data (attrition bias) Treatment failure at 1 month or less	Low risk	
Selective reporting (reporting bias)	Low risk	
Other bias	Unclear risk	Funding not reported

Puczynski 1987

Methods	Randomised controlled, double-blind trial
Participants	Children > 2 years old enrolled May 1984 to February 1985 in Illinois, US
Interventions	Amoxicillin 100 mg x 1 dose followed by placebo 3 times daily for 10 days versus placebo x 1 dose followed by amoxicillin 40 mg/kg 3 times daily for 10 days
Outcomes	Absence of fever and otalgia, decreased irritability and improved appearance of tympanic membrane at 48 to 72 hours and 10 to 14 days after study entry
Notes	-

Risk of bias

Bias	Authors' judgement	Support for judgement
Blinding (performance bias and detection bias) Treatment failure at 1 month or less	Low risk	
Incomplete outcome data (attrition bias) Treatment failure at 1 month or less	Low risk	
Selective reporting (reporting bias)	Low risk	
Other bias	High risk	Stopped early with small sample size due to treatment failures in experimental group, funding not reported

Rodriguez 1996

Methods	RCT	
Participants	Multicentre trial of children 6 months to 12 years old in Guatemala	
Interventions	Azithromycin 10 mg/kg once daily for 3 days versus cefaclor 40 mg/kg 3 times daily for 10 days	
Outcomes	Absence or improvement of signs and symptoms at 10 to 14 days after study entry	
Notes	-	
<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement
Blinding (performance bias and detection bias) Treatment failure at 1 month or less	High risk	
Incomplete outcome data (attrition bias) Treatment failure at 1 month or less	Low risk	
Selective reporting (reporting bias)	Low risk	
Other bias	Unclear risk	Funding not reported

Schaad 1993

Methods	RCT	
Participants	Swiss multicentre trial of children 2 to 12 years old	
Interventions	Azithromycin 10 mg/kg once daily for 3 days versus amoxil-clavulanate 40 mg/kg 3 times daily for 10 days	
Outcomes	Absence or improvement of signs and symptoms at 12 to 16 days after study entry	
Notes	-	
<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement
Blinding (performance bias and detection bias) Treatment failure at 1 month or less	High risk	

Schaad 1993 (Continued)

Incomplete outcome data (attrition bias) Treatment failure at 1 month or less	Low risk	
Selective reporting (reporting bias)	Low risk	
Other bias	Unclear risk	Funding not reported

Varsano 1988

Methods	Randomised controlled, double-blind trial
Participants	Children 6 months to 8 years old seen at a medical clinic in Israel
Interventions	Ceftriaxone 50 mg/kg intramuscularly x 1 dose + placebo 3 times daily for 7 days versus placebo intramuscularly x 1 dose + amoxicillin 12.5 mg/kg 3 times daily for 7 days
Outcomes	Absence of signs and symptoms with or without middle ear effusion at 7 to 10 days and no recurrence at 30 days after study entry
Notes	-

Risk of bias

Bias	Authors' judgement	Support for judgement
Other bias	Unclear risk	Industry funded

Varsano 1997

Methods	RCT
Participants	Israeli children aged 4 months to 6 years
Interventions	Ceftriaxone 50 mg/kg single intramuscular dose versus 37.5/9.4 mg/kg/day amoxicillin/clavulanate for 10 days
Outcomes	Resolution of symptoms and no recurrence at 11 days, no relapse at 30, 60 and 90 days after study entry
Notes	-

Risk of bias

Bias	Authors' judgement	Support for judgement
------	--------------------	-----------------------

Varsano 1997 (Continued)

Random sequence generation (selection bias)	Low risk	
Blinding (performance bias and detection bias) Treatment failure at 1 month or less	High risk	
Incomplete outcome data (attrition bias) Treatment failure at 1 month or less	Low risk	
Selective reporting (reporting bias)	High risk	Separate analysis performed for simple and recurrent AOM. Findings only reported in cases where differences reached statistical significance
Other bias	Unclear risk	Industry funded

Wang 2004

Methods	Prospective comparative randomized trial
Participants	Chinese children between 3 months and 6 years of age
Interventions	Ceftriaxone 50 mg/kg single intramuscular dose versus amoxicillin/clavulanate 45 mg/kg/day 3 times daily for 10 days
Outcomes	Absence or improvement of signs and symptoms at day 11 and day 28 after study entry
Notes	-

Risk of bias

Bias	Authors' judgement	Support for judgement
Blinding (performance bias and detection bias) Treatment failure at 1 month or less	High risk	
Incomplete outcome data (attrition bias) Treatment failure at 1 month or less	Low risk	
Selective reporting (reporting bias)	Low risk	
Other bias	Unclear risk	Funding not reported

d: day; GI: gastrointestinal; RCT: randomized controlled trial

Characteristics of excluded studies *[ordered by study ID]*

Study	Reason for exclusion
Bain 1985	Outcomes not reported as treatment failures
Gehanno 1990	Duration of antibiotic in both treatment arms was less than 7 days
Harrison 1993	Duration of antibiotic therapy equivalent in both arms
Jacobson 1979	Duration of antibiotic therapy equivalent in both arms
Jenner 1987	Duration of antibiotic in both treatment arms was less than 7 days
Johnson 1991	Duration of antibiotic therapy equivalent in both arms
Jones 1986	Outcomes not reported as treatment failures
MacLouglin 1996	Duration of antibiotic therapy equivalent in both arms
Mandel 1995	Duration of antibiotic in both treatment arms 7 days or greater
Murph 1993	Duration of antibiotic therapy equivalent in both arms
O'Doherty 1996	Trial participants predominantly adults
Roos 2000	Antibiotic ceftibuten was deemed to be not clinically relevant
Rubenstein 1965	Antibiotic therapy no longer current
Scott 1990	No comparison of different durations of antibiotic
Simon 1997	Antibiotic ceftibuten was deemed to be not clinically relevant
Spencer 1993	Duration of antibiotic therapy equivalent in both arms
Stickler 1964	Antibiotic therapy no longer current
Stickler 1967	Antibiotic therapy no longer current
Suzuki 2009	Duration of antibiotic therapy equivalent in both arms
Varsano 1994	Insufficient data reported in abstract

DATA AND ANALYSES

Comparison 1. Short-acting antibiotic =< 48 hours in short treatment arm

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Treatment failure at 1 month or less	2	118	Peto Odds Ratio (Peto, Fixed, 95% CI)	2.99 [1.04, 8.54]

Comparison 2. Short-acting antibiotic > 48 hours short-term treatment

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Treatment failure at 1 month or less	16	5093	Peto Odds Ratio (Peto, Fixed, 95% CI)	1.34 [1.15, 1.55]
2 Treatment failure at 8 to 19 days	11	3932	Peto Odds Ratio (Peto, Fixed, 95% CI)	1.37 [1.15, 1.64]
3 Treatment failure at 20 to 30 days	9	2476	Peto Odds Ratio (Peto, Fixed, 95% CI)	1.16 [0.94, 1.42]
4 Treatment failure at 3 months or less	7	2068	Peto Odds Ratio (Peto, Fixed, 95% CI)	1.18 [0.98, 1.41]
5 Treatment failure at 90 days	2	207	Peto Odds Ratio (Peto, Fixed, 95% CI)	1.16 [0.65, 2.06]
6 Treatment failure at 30 to 45 days	5	1861	Peto Odds Ratio (Peto, Fixed, 95% CI)	1.18 [0.97, 1.43]

Comparison 3. Short-acting antibiotic > 48 hours short-term treatment, < 2 yrs old

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Treatment failure at 1 month or less	5	570	Peto Odds Ratio (Peto, Fixed, 95% CI)	1.09 [0.76, 1.57]

Comparison 4. Short-acting antibiotic > 48 hours short-term treatment, => 2 yrs old

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Treatment failure at 1 month or less	6	1064	Peto Odds Ratio (Peto, Fixed, 95% CI)	0.85 [0.60, 1.21]

Comparison 5. Short-acting antibiotic > 48 hours short-term treatment, perforated eardrum

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Treatment failure at 1 month or less	1	27	Peto Odds Ratio (Peto, Fixed, 95% CI)	3.62 [0.81, 16.06]

Comparison 6. Short-acting antibiotic > 48 hours short-term treatment, non-perforated eardrum

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Treatment failure at 1 month or less	1	101	Peto Odds Ratio (Peto, Fixed, 95% CI)	1.06 [0.40, 2.75]

Comparison 7. Short-acting antibiotic > 48 hours short-term treatment; Sensitivity Analysis: include chronic OM

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Treatment failure at 1 month or less	6	1713	Peto Odds Ratio (Peto, Fixed, 95% CI)	1.37 [1.09, 1.72]
2 Treatment failure at 20 to 30 days	5	1149	Peto Odds Ratio (Peto, Fixed, 95% CI)	1.17 [0.89, 1.54]

Comparison 8. Short-acting antibiotic > 48 hours short-term treatment; Sensitivity Analysis: exclude chronic OM

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Treatment failure at 1 month or less	4	1298	Peto Odds Ratio (Peto, Fixed, 95% CI)	1.17 [0.73, 1.85]
2 Treatment failure at 20 to 30 days	2	656	Peto Odds Ratio (Peto, Fixed, 95% CI)	1.55 [0.79, 3.04]

Comparison 9. Short-acting antibiotic > 48 hours short-term treatment; Sensitivity analysis: outcome only if "cured"

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Treatment failure at 1 month or less	9	2955	Peto Odds Ratio (Peto, Fixed, 95% CI)	1.31 [1.09, 1.58]
2 Treatment failure at 20 to 30 days	6	1749	Peto Odds Ratio (Peto, Fixed, 95% CI)	1.25 [0.98, 1.58]

Comparison 10. Short-acting antibiotic > 48 hours short-term treatment; Sensitivity analysis: same antibiotic in treatment arms

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Treatment failure at 1 month or less	9	3321	Peto Odds Ratio (Peto, Fixed, 95% CI)	1.65 [1.35, 2.01]
2 Treatment failure at 8 to 19 days	6	2153	Peto Odds Ratio (Peto, Fixed, 95% CI)	1.97 [1.54, 2.52]
3 Treatment failure at 20 to 30 days	4	1319	Peto Odds Ratio (Peto, Fixed, 95% CI)	1.27 [0.92, 1.76]
4 Treatment failure at 3 months or less	5	1492	Peto Odds Ratio (Peto, Fixed, 95% CI)	1.24 [1.00, 1.53]
5 Treatment failure at 90 days	2	207	Peto Odds Ratio (Peto, Fixed, 95% CI)	1.16 [0.65, 2.06]
6 Treatment failure at 30 to 45 days	3	1285	Peto Odds Ratio (Peto, Fixed, 95% CI)	1.25 [1.00, 1.57]

Comparison 11. Short-acting antibiotic > 48 hours short-term treatment, excluding amoxicillin-clavulanate

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Gastrointestinal adverse effects	5	1088	Peto Odds Ratio (Peto, Fixed, 95% CI)	1.06 [0.61, 1.85]

Comparison 12. Short-acting antibiotic > 48 hours short-term treatment; Sensitivity analysis: treatment failure at 1 month or less and risk of bias

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Sensitivity analysis: allocation concealment unclear risk of bias	13	3953	Peto Odds Ratio (Peto, Fixed, 95% CI)	1.30 [1.09, 1.54]
2 Sensitivity analysis: allocation concealment low risk of bias	3	1140	Peto Odds Ratio (Peto, Fixed, 95% CI)	1.45 [1.08, 1.93]
3 Sensitivity analysis: blinding high and unclear risk of bias	12	3927	Peto Odds Ratio (Peto, Fixed, 95% CI)	1.18 [1.00, 1.40]
4 Sensitivity analysis: blinding low risk of bias	4	1166	Peto Odds Ratio (Peto, Fixed, 95% CI)	2.03 [1.48, 2.77]

Comparison 13. Short-acting antibiotic > 48 hours short-term treatment; Sensitivity analysis amoxil-clav. both arms

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Amoxil-clav. 5 versus 10 days, treatment failure 1 month or less	2	942	Peto Odds Ratio (Peto, Fixed, 95% CI)	1.99 [1.44, 2.74]
2 Excluding amoxil-clav. 5 versus 10 days, treatment failure 1 month or less	14	4151	Peto Odds Ratio (Peto, Fixed, 95% CI)	1.20 [1.02, 1.42]

Comparison 14. Short-acting antibiotic > 48 hours short-term treatment, => 2 yrs old; Sensitivity analysis

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Treatment failure at 1 month or less	5	656	Peto Odds Ratio (Peto, Fixed, 95% CI)	1.00 [0.68, 1.47]

Comparison 15. Short-acting antibiotic > 48 hours short-term treatment, < 2 yrs old; Sensitivity analysis

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Treatment failure at 1 month or less	5	548	Peto Odds Ratio (Peto, Fixed, 95% CI)	1.19 [0.82, 1.73]

Comparison 16. Short-acting antibiotic, > 48 hours short-term treatment, adverse GI effects

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Gastrointestinal adverse effects	13	4918	Peto Odds Ratio (Peto, Fixed, 95% CI)	0.72 [0.60, 0.87]

Comparison 17. Ceftriaxone

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Treatment failure at 1 month or less	8	1709	Peto Odds Ratio (Peto, Fixed, 95% CI)	1.07 [0.86, 1.33]
2 Treatment failure at 3 months or less	3	701	Peto Odds Ratio (Peto, Fixed, 95% CI)	0.89 [0.66, 1.21]
3 Gastrointestinal adverse effects	1	402	Peto Odds Ratio (Peto, Fixed, 95% CI)	2.89 [1.70, 4.91]

Comparison 18. Azithromycin single-dose short-term treatment

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Treatment failure at 25 to 32 days	2	608	Peto Odds Ratio (Peto, Fixed, 95% CI)	1.01 [0.69, 1.47]
2 Treatment failure at 25 to 32 days, =< 2 yrs old	2	368	Peto Odds Ratio (Peto, Fixed, 95% CI)	0.68 [0.43, 1.09]
3 Treatment failure at 25 to 32 days, > 2 yrs old	2	240	Peto Odds Ratio (Peto, Fixed, 95% CI)	1.08 [0.53, 2.20]
4 Gastrointestinal adverse effects	2	658	Peto Odds Ratio (Peto, Fixed, 95% CI)	0.66 [0.45, 0.96]

Comparison 19. Azithromycin 3 to 5 days short-term treatment

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Treatment failure at 1 month or less	19	4354	Peto Odds Ratio (Peto, Fixed, 95% CI)	1.02 [0.87, 1.20]
2 Treatment failure at 8 to 19 days	18	4347	Peto Odds Ratio (Peto, Fixed, 95% CI)	1.27 [1.04, 1.55]
3 Treatment failure at 20 to 30 days	11	2708	Peto Odds Ratio (Peto, Fixed, 95% CI)	0.98 [0.82, 1.17]

Comparison 20. Azithromycin 3 to 5 days short-term treatment, < 2 years old

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Treatment failure at 1 month or less	2	138	Peto Odds Ratio (Peto, Fixed, 95% CI)	1.92 [0.73, 5.04]

Comparison 21. Azithromycin 3 to 5 days short-term treatment, => 2 years old

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Treatment failure at 1 month or less	2	656	Peto Odds Ratio (Peto, Fixed, 95% CI)	1.34 [0.61, 2.94]

Comparison 22. Azithromycin 3-5 days short-term treatment; Sensitivity analysis: include chronic OM

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Treatment failure at 1 month or less	7	1688	Peto Odds Ratio (Peto, Fixed, 95% CI)	0.96 [0.70, 1.31]
2 Treatment failure at 20 to 30 days	4	740	Peto Odds Ratio (Peto, Fixed, 95% CI)	0.83 [0.57, 1.21]

Comparison 23. Azithromycin 3 to 5 days short-term treatment; Sensitivity analysis: exclude chronic OM

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Treatment failure at 1 month or less	6	1193	Peto Odds Ratio (Peto, Fixed, 95% CI)	1.26 [0.89, 1.80]
2 Treatment failure at 20 to 30 days	3	688	Peto Odds Ratio (Peto, Fixed, 95% CI)	1.27 [0.86, 1.86]

Comparison 24. Azithromycin 3 to 5 days short-term treatment; Sensitivity analysis: outcome only if “cured”

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Treatment failure at 1 month or less	11	2355	Peto Odds Ratio (Peto, Fixed, 95% CI)	0.74 [0.61, 0.90]
2 Treatment failure at 20 to 30 days	5	902	Peto Odds Ratio (Peto, Fixed, 95% CI)	0.96 [0.70, 1.31]

Comparison 25. Azithromycin 3 to 5 days short-term treatment; Sensitivity analysis: similar spectrum antibiotic in treatment arms

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Treatment failure at 20 to 30 days	6	1254	Peto Odds Ratio (Peto, Fixed, 95% CI)	1.02 [0.78, 1.34]
2 Treatment failure at 1 month or less	9	2205	Peto Odds Ratio (Peto, Fixed, 95% CI)	1.13 [0.89, 1.44]

Comparison 26. Azithromycin 3 to 5 days short-term treatment; Sensitivity analysis: risk of bias

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Sensitivity analysis: allocation concealment unclear or high risk of bias	17	4165	Odds Ratio (M-H, Fixed, 95% CI)	1.04 [0.88, 1.22]
2 Sensitivity analysis: allocation concealment low risk of bias	2	189	Peto Odds Ratio (Peto, Fixed, 95% CI)	0.13 [0.02, 0.90]
3 Sensitivity analysis: blinding high and unclear risk of bias	15	3348	Peto Odds Ratio (Peto, Fixed, 95% CI)	1.15 [0.94, 1.40]
4 Sensitivity analysis: blinding low risk of bias	4	1006	Peto Odds Ratio (Peto, Fixed, 95% CI)	0.80 [0.60, 1.07]

Comparison 27. Azithromycin 3 to 5 days short-term treatment; Sensitivity analysis: industry funding

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Treatment failure 1 month or less	8	1747	Peto Odds Ratio (Peto, Fixed, 95% CI)	0.97 [0.78, 1.21]
2 Treatment failure at 8 to 19 days	7	1667	Peto Odds Ratio (Peto, Fixed, 95% CI)	1.43 [1.10, 1.87]
3 Treatment failure at 20 to 30 days	6	1469	Peto Odds Ratio (Peto, Fixed, 95% CI)	0.93 [0.74, 1.17]

Comparison 28. Azithromycin 3 to 5 days short-term treatment; Sensitivity analysis: funding not reported

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Treatment failure 1 month or less	11	2616	Peto Odds Ratio (Peto, Fixed, 95% CI)	1.09 [0.86, 1.38]
2 Treatment failure at 8 to 19 days	11	2680	Peto Odds Ratio (Peto, Fixed, 95% CI)	1.08 [0.80, 1.46]
3 Treatment failure at 20 to 30 days	5	1239	Peto Odds Ratio (Peto, Fixed, 95% CI)	1.06 [0.81, 1.39]

Comparison 29. Azithromycin 3 to 5 days short-term treatment, adverse GI effects

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Gastrointestinal adverse effects	14	3722	Peto Odds Ratio (Peto, Fixed, 95% CI)	0.36 [0.28, 0.46]

Comparison 30. Methods - combined control group versus twice counting tx

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Twice counting - short-acting antibiotic > 48 hours short-term treatment, treatment failure at 1 month or less	9	2249	Peto Odds Ratio (Peto, Fixed, 95% CI)	1.43 [1.14, 1.79]
2 Combined control groups - short-acting antibiotic > 48 hours short-term treatment, treatment failure 1 month or less	9	2611	Peto Odds Ratio (Peto, Fixed, 95% CI)	1.36 [1.10, 1.67]
3 Twice counting - short-acting antibiotic > 48 hours short-term treatment, treatment failure at 1 to 19 days	4	1138	Peto Odds Ratio (Peto, Fixed, 95% CI)	1.52 [1.11, 2.10]
4 Combined control groups - short-acting antibiotic > 48 hours short-term treatment, treatment failure 1 to 19 days	4	1327	Peto Odds Ratio (Peto, Fixed, 95% CI)	1.45 [1.08, 1.95]
5 Twice counting - short-acting antibiotic > 48 hours short-term treatment, treatment failure 20 to 30 days	7	1632	Peto Odds Ratio (Peto, Fixed, 95% CI)	1.26 [0.97, 1.64]
6 Combined control groups - short-acting antibiotic > 48 hours short-term treatment, treatment failure at 20 to 30 days	7	1805	Peto Odds Ratio (Peto, Fixed, 95% CI)	1.22 [0.95, 1.57]
7 Twice counting - short-acting antibiotic > 48 hours short-term treatment, treatment failure at 30 to 40 days	2	538	Peto Odds Ratio (Peto, Fixed, 95% CI)	1.07 [0.74, 1.55]
8 Combined control groups - short-acting antibiotic > 48 hours short-term treatment, treatment failure 30 to 40 days	2	686	Peto Odds Ratio (Peto, Fixed, 95% CI)	1.12 [0.80, 1.57]


9 Twice counting - short-acting antibiotic > 48 hours short-term treatment, treatment failure at 3 months or less	4	745	Peto Odds Ratio (Peto, Fixed, 95% CI)	1.10 [0.80, 1.50]
10 Combined control groups - short-acting antibiotic > 48 hours short-term treatment, treatment failure at 3 months or less	4	893	Peto Odds Ratio (Peto, Fixed, 95% CI)	1.13 [0.85, 1.51]

Analysis 1.1. Comparison 1 Short-acting antibiotic ≤ 48 hours in short treatment arm, Outcome 1 Treatment failure at 1 month or less.

Review: Short-course antibiotics for acute otitis media

Comparison: 1 Short-acting antibiotic ≤ 48 hours in short treatment arm

Outcome: 1 Treatment failure at 1 month or less




Analysis 2.1. Comparison 2 Short-acting antibiotic > 48 hours short-term treatment, Outcome 1 Treatment failure at 1 month or less.

Review: Short-course antibiotics for acute otitis media

Comparison: 2 Short-acting antibiotic > 48 hours short-term treatment

Outcome: 1 Treatment failure at 1 month or less




Analysis 2.2. Comparison 2 Short-acting antibiotic > 48 hours short-term treatment, Outcome 2 Treatment failure at 8 to 19 days.

Review: Short-course antibiotics for acute otitis media

Comparison: 2 Short-acting antibiotic > 48 hours short-term treatment

Outcome: 2 Treatment failure at 8 to 19 days




Analysis 2.3. Comparison 2 Short-acting antibiotic > 48 hours short-term treatment, Outcome 3 Treatment failure at 20 to 30 days.

Review: Short-course antibiotics for acute otitis media

Comparison: 2 Short-acting antibiotic > 48 hours short-term treatment

Outcome: 3 Treatment failure at 20 to 30 days




Analysis 2.4. Comparison 2 Short-acting antibiotic > 48 hours short-term treatment, Outcome 4 Treatment failure at 3 months or less.

Review: Short-course antibiotics for acute otitis media

Comparison: 2 Short-acting antibiotic > 48 hours short-term treatment

Outcome: 4 Treatment failure at 3 months or less




Analysis 2.5. Comparison 2 Short-acting antibiotic > 48 hours short-term treatment, Outcome 5 Treatment failure at 90 days.

Review: Short-course antibiotics for acute otitis media

Comparison: 2 Short-acting antibiotic > 48 hours short-term treatment

Outcome: 5 Treatment failure at 90 days




Analysis 2.6. Comparison 2 Short-acting antibiotic > 48 hours short-term treatment, Outcome 6 Treatment failure at 30 to 45 days.

Review: Short-course antibiotics for acute otitis media

Comparison: 2 Short-acting antibiotic > 48 hours short-term treatment

Outcome: 6 Treatment failure at 30 to 45 days




Analysis 3.1. Comparison 3 Short-acting antibiotic > 48 hours short-term treatment, < 2 yrs old, Outcome 1 Treatment failure at 1 month or less.

Review: Short-course antibiotics for acute otitis media

Comparison: 3 Short-acting antibiotic > 48 hours short-term treatment, < 2 yrs old

Outcome: 1 Treatment failure at 1 month or less




Analysis 4.1. Comparison 4 Short-acting antibiotic > 48 hours short-term treatment, => 2 yrs old, Outcome 1 Treatment failure at 1 month or less.

Review: Short-course antibiotics for acute otitis media

Comparison: 4 Short-acting antibiotic > 48 hours short-term treatment, => 2 yrs old

Outcome: 1 Treatment failure at 1 month or less




Analysis 5.1. Comparison 5 Short-acting antibiotic > 48 hours short-term treatment, perforated eardrum, Outcome 1 Treatment failure at 1 month or less.

Review: Short-course antibiotics for acute otitis media

Comparison: 5 Short-acting antibiotic > 48 hours short-term treatment, perforated eardrum

Outcome: 1 Treatment failure at 1 month or less




Analysis 6.1. Comparison 6 Short-acting antibiotic > 48 hours short-term treatment, non-perforated eardrum, Outcome 1 Treatment failure at 1 month or less.

Review: Short-course antibiotics for acute otitis media

Comparison: 6 Short-acting antibiotic > 48 hours short-term treatment, non-perforated eardrum

Outcome: 1 Treatment failure at 1 month or less




Analysis 7.1. Comparison 7 Short-acting antibiotic > 48 hours short-term treatment; Sensitivity Analysis: include chronic OM, Outcome 1 Treatment failure at 1 month or less.

Review: Short-course antibiotics for acute otitis media

Comparison: 7 Short-acting antibiotic > 48 hours short-term treatment; Sensitivity Analysis: include chronic OM

Outcome: 1 Treatment failure at 1 month or less




Analysis 7.2. Comparison 7 Short-acting antibiotic > 48 hours short-term treatment; Sensitivity Analysis: include chronic OM, Outcome 2 Treatment failure at 20 to 30 days.

Review: Short-course antibiotics for acute otitis media

Comparison: 7 Short-acting antibiotic > 48 hours short-term treatment; Sensitivity Analysis: include chronic OM

Outcome: 2 Treatment failure at 20 to 30 days




Analysis 8.1. Comparison 8 Short-acting antibiotic > 48 hours short-term treatment; Sensitivity Analysis: exclude chronic OM, Outcome 1 Treatment failure at 1 month or less.

Review: Short-course antibiotics for acute otitis media

Comparison: 8 Short-acting antibiotic > 48 hours short-term treatment; Sensitivity Analysis: exclude chronic OM

Outcome: 1 Treatment failure at 1 month or less




Analysis 8.2. Comparison 8 Short-acting antibiotic > 48 hours short-term treatment; Sensitivity Analysis: exclude chronic OM, Outcome 2 Treatment failure at 20 to 30 days.

Review: Short-course antibiotics for acute otitis media

Comparison: 8 Short-acting antibiotic > 48 hours short-term treatment; Sensitivity Analysis: exclude chronic OM

Outcome: 2 Treatment failure at 20 to 30 days




Analysis 9.1. Comparison 9 Short-acting antibiotic > 48 hours short-term treatment; Sensitivity analysis: outcome only if “cured”, Outcome 1 Treatment failure at 1 month or less.

Review: Short-course antibiotics for acute otitis media

Comparison: 9 Short-acting antibiotic > 48 hours short-term treatment; Sensitivity analysis: outcome only if “cured”

Outcome: 1 Treatment failure at 1 month or less




Analysis 9.2. Comparison 9 Short-acting antibiotic > 48 hours short-term treatment; Sensitivity analysis: outcome only if "cured", Outcome 2 Treatment failure at 20 to 30 days.

Review: Short-course antibiotics for acute otitis media

Comparison: 9 Short-acting antibiotic > 48 hours short-term treatment; Sensitivity analysis: outcome only if "cured"

Outcome: 2 Treatment failure at 20 to 30 days




Analysis 10.1. Comparison 10 Short-acting antibiotic > 48 hours short-term treatment; Sensitivity analysis: same antibiotic in treatment arms, Outcome 1 Treatment failure at 1 month or less.

Review: Short-course antibiotics for acute otitis media

Comparison: 10 Short-acting antibiotic > 48 hours short-term treatment; Sensitivity analysis: same antibiotic in treatment arms

Outcome: 1 Treatment failure at 1 month or less




Analysis 10.2. Comparison 10 Short-acting antibiotic > 48 hours short-term treatment; Sensitivity analysis: same antibiotic in treatment arms, Outcome 2 Treatment failure at 8 to 19 days.

Review: Short-course antibiotics for acute otitis media

Comparison: 10 Short-acting antibiotic > 48 hours short-term treatment; Sensitivity analysis: same antibiotic in treatment arms

Outcome: 2 Treatment failure at 8 to 19 days




Analysis 10.3. Comparison 10 Short-acting antibiotic > 48 hours short-term treatment; Sensitivity analysis: same antibiotic in treatment arms, Outcome 3 Treatment failure at 20 to 30 days.

Review: Short-course antibiotics for acute otitis media

Comparison: 10 Short-acting antibiotic > 48 hours short-term treatment; Sensitivity analysis: same antibiotic in treatment arms

Outcome: 3 Treatment failure at 20 to 30 days




Analysis 10.4. Comparison 10 Short-acting antibiotic > 48 hours short-term treatment; Sensitivity analysis: same antibiotic in treatment arms, Outcome 4 Treatment failure at 3 months or less.

Review: Short-course antibiotics for acute otitis media

Comparison: 10 Short-acting antibiotic > 48 hours short-term treatment; Sensitivity analysis: same antibiotic in treatment arms

Outcome: 4 Treatment failure at 3 months or less




Analysis 10.5. Comparison 10 Short-acting antibiotic > 48 hours short-term treatment; Sensitivity analysis: same antibiotic in treatment arms, Outcome 5 Treatment failure at 90 days.

Review: Short-course antibiotics for acute otitis media

Comparison: 10 Short-acting antibiotic > 48 hours short-term treatment; Sensitivity analysis: same antibiotic in treatment arms

Outcome: 5 Treatment failure at 90 days




Analysis 10.6. Comparison 10 Short-acting antibiotic > 48 hours short-term treatment; Sensitivity analysis: same antibiotic in treatment arms, Outcome 6 Treatment failure at 30 to 45 days.

Review: Short-course antibiotics for acute otitis media

Comparison: 10 Short-acting antibiotic > 48 hours short-term treatment; Sensitivity analysis: same antibiotic in treatment arms

Outcome: 6 Treatment failure at 30 to 45 days




Analysis 11.1. Comparison 11 Short-acting antibiotic > 48 hours short-term treatment, excluding amoxicillin-clavulanate, Outcome 1 Gastrointestinal adverse effects.

Review: Short-course antibiotics for acute otitis media

Comparison: 11 Short-acting antibiotic > 48 hours short-term treatment, excluding amoxicillin-clavulanate

Outcome: 1 Gastrointestinal adverse effects




Analysis 12.1. Comparison 12 Short-acting antibiotic > 48 hours short-term treatment; Sensitivity analysis: treatment failure at 1 month or less and risk of bias, Outcome 1 Sensitivity analysis: allocation concealment unclear risk of bias.

Review: Short-course antibiotics for acute otitis media

Comparison: 12 Short-acting antibiotic > 48 hours short-term treatment; Sensitivity analysis: treatment failure at 1 month or less and risk of bias

Outcome: 1 Sensitivity analysis: allocation concealment unclear risk of bias




Analysis 12.2. Comparison 12 Short-acting antibiotic > 48 hours short-term treatment; Sensitivity analysis: treatment failure at 1 month or less and risk of bias, Outcome 2 Sensitivity analysis: allocation concealment low risk of bias.

Review: Short-course antibiotics for acute otitis media

Comparison: 12 Short-acting antibiotic > 48 hours short-term treatment; Sensitivity analysis: treatment failure at 1 month or less and risk of bias

Outcome: 2 Sensitivity analysis: allocation concealment low risk of bias




Analysis 12.3. Comparison 12 Short-acting antibiotic > 48 hours short-term treatment; Sensitivity analysis: treatment failure at 1 month or less and risk of bias, Outcome 3 Sensitivity analysis: blinding high and unclear risk of bias.

Review: Short-course antibiotics for acute otitis media

Comparison: 12 Short-acting antibiotic > 48 hours short-term treatment; Sensitivity analysis: treatment failure at 1 month or less and risk of bias

Outcome: 3 Sensitivity analysis: blinding high and unclear risk of bias




Analysis 12.4. Comparison 12 Short-acting antibiotic > 48 hours short-term treatment; Sensitivity analysis: treatment failure at 1 month or less and risk of bias, Outcome 4 Sensitivity analysis: blinding low risk of bias.

Review: Short-course antibiotics for acute otitis media

Comparison: 12 Short-acting antibiotic > 48 hours short-term treatment; Sensitivity analysis: treatment failure at 1 month or less and risk of bias

Outcome: 4 Sensitivity analysis: blinding low risk of bias




Analysis 13.1. Comparison 13 Short-acting antibiotic > 48 hours short-term treatment; Sensitivity analysis amoxil-clav. both arms, Outcome 1 Amoxil-clav. 5 versus 10 days, treatment failure 1 month or less.

Review: Short-course antibiotics for acute otitis media

Comparison: 13 Short-acting antibiotic > 48 hours short-term treatment; Sensitivity analysis amoxil-clav. both arms

Outcome: 1 Amoxil-clav. 5 versus 10 days, treatment failure 1 month or less




Analysis 13.2. Comparison 13 Short-acting antibiotic > 48 hours short-term treatment; Sensitivity analysis amoxil-clav. both arms, Outcome 2 Excluding amoxil-clav. 5 versus 10 days, treatment failure 1 month or less.

Review: Short-course antibiotics for acute otitis media

Comparison: 13 Short-acting antibiotic > 48 hours short-term treatment; Sensitivity analysis amoxil-clav. both arms

Outcome: 2 Excluding amoxil-clav. 5 versus 10 days, treatment failure 1 month or less




Analysis 14.1. Comparison 14 Short-acting antibiotic > 48 hours short-term treatment, => 2 yrs old; Sensitivity analysis, Outcome 1 Treatment failure at 1 month or less.

Review: Short-course antibiotics for acute otitis media

Comparison: 14 Short-acting antibiotic > 48 hours short-term treatment, => 2 yrs old; Sensitivity analysis

Outcome: 1 Treatment failure at 1 month or less




Analysis 15.1. Comparison 15 Short-acting antibiotic > 48 hours short-term treatment, < 2 yrs old; Sensitivity analysis, Outcome 1 Treatment failure at 1 month or less.

Review: Short-course antibiotics for acute otitis media

Comparison: 15 Short-acting antibiotic > 48 hours short-term treatment, < 2 yrs old; Sensitivity analysis

Outcome: 1 Treatment failure at 1 month or less




Analysis 16.1. Comparison 16 Short-acting antibiotic, > 48 hours short-term treatment, adverse GI effects, Outcome 1 Gastrointestinal adverse effects.

Review: Short-course antibiotics for acute otitis media

Comparison: 16 Short-acting antibiotic, > 48 hours short-term treatment, adverse GI effects

Outcome: 1 Gastrointestinal adverse effects




Analysis 17.1. Comparison 17 Ceftriaxone, Outcome 1 Treatment failure at 1 month or less.

Review: Short-course antibiotics for acute otitis media

Comparison: 17 Ceftriaxone

Outcome: 1 Treatment failure at 1 month or less




Analysis 17.2. Comparison 17 Ceftriaxone, Outcome 2 Treatment failure at 3 months or less.

Review: Short-course antibiotics for acute otitis media

Comparison: 17 Ceftriaxone

Outcome: 2 Treatment failure at 3 months or less




Analysis 17.3. Comparison 17 Ceftriaxone, Outcome 3 Gastrointestinal adverse effects.

Review: Short-course antibiotics for acute otitis media

Comparison: 17 Ceftriaxone

Outcome: 3 Gastrointestinal adverse effects




Analysis 18.1. Comparison 18 Azithromycin single-dose short-term treatment, Outcome 1 Treatment failure at 25 to 32 days.

Review: Short-course antibiotics for acute otitis media

Comparison: 18 Azithromycin single-dose short-term treatment

Outcome: 1 Treatment failure at 25 to 32 days




Analysis 18.2. Comparison 18 Azithromycin single-dose short-term treatment, Outcome 2 Treatment failure at 25 to 32 days, =< 2 yrs old.

Review: Short-course antibiotics for acute otitis media

Comparison: 18 Azithromycin single-dose short-term treatment

Outcome: 2 Treatment failure at 25 to 32 days, =< 2 yrs old




Analysis 18.3. Comparison 18 Azithromycin single-dose short-term treatment, Outcome 3 Treatment failure at 25 to 32 days, > 2 yrs old.

Review: Short-course antibiotics for acute otitis media

Comparison: 18 Azithromycin single-dose short-term treatment

Outcome: 3 Treatment failure at 25 to 32 days, > 2 yrs old




Analysis 18.4. Comparison 18 Azithromycin single-dose short-term treatment, Outcome 4 Gastrointestinal adverse effects.

Review: Short-course antibiotics for acute otitis media

Comparison: 18 Azithromycin single-dose short-term treatment

Outcome: 4 Gastrointestinal adverse effects




Analysis 19.1. Comparison 19 Azithromycin 3 to 5 days short-term treatment, Outcome 1 Treatment failure at 1 month or less.

Review: Short-course antibiotics for acute otitis media

Comparison: 19 Azithromycin 3 to 5 days short-term treatment

Outcome: 1 Treatment failure at 1 month or less




Analysis 19.2. Comparison 19 Azithromycin 3 to 5 days short-term treatment, Outcome 2 Treatment failure at 8 to 19 days.

Review: Short-course antibiotics for acute otitis media

Comparison: 19 Azithromycin 3 to 5 days short-term treatment

Outcome: 2 Treatment failure at 8 to 19 days




Analysis 19.3. Comparison 19 Azithromycin 3 to 5 days short-term treatment, Outcome 3 Treatment failure at 20 to 30 days.

Review: Short-course antibiotics for acute otitis media

Comparison: 19 Azithromycin 3 to 5 days short-term treatment

Outcome: 3 Treatment failure at 20 to 30 days




Analysis 20.1. Comparison 20 Azithromycin 3 to 5 days short-term treatment, < 2 years old, Outcome 1 Treatment failure at 1 month or less.

Review: Short-course antibiotics for acute otitis media

Comparison: 20 Azithromycin 3 to 5 days short-term treatment, < 2 years old

Outcome: 1 Treatment failure at 1 month or less




Analysis 21.1. Comparison 21 Azithromycin 3 to 5 days short-term treatment, => 2 years old, Outcome 1 Treatment failure at 1 month or less.

Review: Short-course antibiotics for acute otitis media

Comparison: 21 Azithromycin 3 to 5 days short-term treatment, => 2 years old

Outcome: 1 Treatment failure at 1 month or less




Analysis 22.1. Comparison 22 Azithromycin 3-5 days short-term treatment; Sensitivity analysis: include chronic OM, Outcome 1 Treatment failure at 1 month or less.

Review: Short-course antibiotics for acute otitis media

Comparison: 22 Azithromycin 3-5 days short-term treatment; Sensitivity analysis: include chronic OM

Outcome: 1 Treatment failure at 1 month or less




Analysis 22.2. Comparison 22 Azithromycin 3-5 days short-term treatment; Sensitivity analysis: include chronic OM, Outcome 2 Treatment failure at 20 to 30 days.

Review: Short-course antibiotics for acute otitis media

Comparison: 22 Azithromycin 3-5 days short-term treatment; Sensitivity analysis: include chronic OM

Outcome: 2 Treatment failure at 20 to 30 days




Analysis 23.1. Comparison 23 Azithromycin 3 to 5 days short-term treatment; Sensitivity analysis: exclude chronic OM, Outcome 1 Treatment failure at 1 month or less.

Review: Short-course antibiotics for acute otitis media

Comparison: 23 Azithromycin 3 to 5 days short-term treatment; Sensitivity analysis: exclude chronic OM

Outcome: 1 Treatment failure at 1 month or less




Analysis 23.2. Comparison 23 Azithromycin 3 to 5 days short-term treatment; Sensitivity analysis: exclude chronic OM, Outcome 2 Treatment failure at 20 to 30 days.

Review: Short-course antibiotics for acute otitis media

Comparison: 23 Azithromycin 3 to 5 days short-term treatment; Sensitivity analysis: exclude chronic OM

Outcome: 2 Treatment failure at 20 to 30 days




Analysis 24.1. Comparison 24 Azithromycin 3 to 5 days short-term treatment; Sensitivity analysis: outcome only if "cured", Outcome 1 Treatment failure at 1 month or less.

Review: Short-course antibiotics for acute otitis media

Comparison: 24 Azithromycin 3 to 5 days short-term treatment; Sensitivity analysis: outcome only if "cured"

Outcome: 1 Treatment failure at 1 month or less




Analysis 24.2. Comparison 24 Azithromycin 3 to 5 days short-term treatment; Sensitivity analysis: outcome only if “cured”, Outcome 2 Treatment failure at 20 to 30 days.

Review: Short-course antibiotics for acute otitis media

Comparison: 24 Azithromycin 3 to 5 days short-term treatment; Sensitivity analysis: outcome only if “cured”

Outcome: 2 Treatment failure at 20 to 30 days




Analysis 25.1. Comparison 25 Azithromycin 3 to 5 days short-term treatment; Sensitivity analysis: similar spectrum antibiotic in treatment arms, Outcome 1 Treatment failure at 20 to 30 days.

Review: Short-course antibiotics for acute otitis media

Comparison: 25 Azithromycin 3 to 5 days short-term treatment; Sensitivity analysis: similar spectrum antibiotic in treatment arms

Outcome: 1 Treatment failure at 20 to 30 days




Analysis 25.2. Comparison 25 Azithromycin 3 to 5 days short-term treatment; Sensitivity analysis: similar spectrum antibiotic in treatment arms, Outcome 2 Treatment failure at 1 month or less.

Review: Short-course antibiotics for acute otitis media

Comparison: 25 Azithromycin 3 to 5 days short-term treatment; Sensitivity analysis: similar spectrum antibiotic in treatment arms

Outcome: 2 Treatment failure at 1 month or less




Analysis 26.1. Comparison 26 Azithromycin 3 to 5 days short-term treatment; Sensitivity analysis: risk of bias, Outcome 1 Sensitivity analysis: allocation concealment unclear or high risk of bias.

Review: Short-course antibiotics for acute otitis media

Comparison: 26 Azithromycin 3 to 5 days short-term treatment; Sensitivity analysis: risk of bias

Outcome: 1 Sensitivity analysis: allocation concealment unclear or high risk of bias




Analysis 26.2. Comparison 26 Azithromycin 3 to 5 days short-term treatment; Sensitivity analysis: risk of bias, Outcome 2 Sensitivity analysis: allocation concealment low risk of bias.

Review: Short-course antibiotics for acute otitis media

Comparison: 26 Azithromycin 3 to 5 days short-term treatment; Sensitivity analysis: risk of bias

Outcome: 2 Sensitivity analysis: allocation concealment low risk of bias




Analysis 26.3. Comparison 26 Azithromycin 3 to 5 days short-term treatment; Sensitivity analysis: risk of bias, Outcome 3 Sensitivity analysis: blinding high and unclear risk of bias.

Review: Short-course antibiotics for acute otitis media

Comparison: 26 Azithromycin 3 to 5 days short-term treatment; Sensitivity analysis: risk of bias

Outcome: 3 Sensitivity analysis: blinding high and unclear risk of bias




Analysis 26.4. Comparison 26 Azithromycin 3 to 5 days short-term treatment; Sensitivity analysis: risk of bias, Outcome 4 Sensitivity analysis: blinding low risk of bias.

Review: Short-course antibiotics for acute otitis media

Comparison: 26 Azithromycin 3 to 5 days short-term treatment; Sensitivity analysis: risk of bias

Outcome: 4 Sensitivity analysis: blinding low risk of bias




Analysis 27.1. Comparison 27 Azithromycin 3 to 5 days short-term treatment; Sensitivity analysis: industry funding, Outcome 1 Treatment failure 1 month or less.

Review: Short-course antibiotics for acute otitis media

Comparison: 27 Azithromycin 3 to 5 days short-term treatment; Sensitivity analysis: industry funding

Outcome: 1 Treatment failure 1 month or less




Analysis 27.2. Comparison 27 Azithromycin 3 to 5 days short-term treatment; Sensitivity analysis: industry funding, Outcome 2 Treatment failure at 8 to 19.

Review: Short-course antibiotics for acute otitis media

Comparison: 27 Azithromycin 3 to 5 days short-term treatment; Sensitivity analysis: industry funding

Outcome: 2 Treatment failure at 8 to 19




Analysis 27.3. Comparison 27 Azithromycin 3 to 5 days short-term treatment; Sensitivity analysis: industry funding, Outcome 3 Treatment failure at 20 to 30 days.

Review: Short-course antibiotics for acute otitis media

Comparison: 27 Azithromycin 3 to 5 days short-term treatment; Sensitivity analysis: industry funding

Outcome: 3 Treatment failure at 20 to 30 days




Analysis 28.1. Comparison 28 Azithromycin 3 to 5 days short-term treatment; Sensitivity analysis: funding not reported, Outcome 1 Treatment failure 1 month or less.

Review: Short-course antibiotics for acute otitis media

Comparison: 28 Azithromycin 3 to 5 days short-term treatment; Sensitivity analysis: funding not reported

Outcome: 1 Treatment failure 1 month or less




Analysis 28.2. Comparison 28 Azithromycin 3 to 5 days short-term treatment; Sensitivity analysis: funding not reported, Outcome 2 Treatment failure at 8 to 19 days.

Review: Short-course antibiotics for acute otitis media

Comparison: 28 Azithromycin 3 to 5 days short-term treatment; Sensitivity analysis: funding not reported

Outcome: 2 Treatment failure at 8 to 19 days




Analysis 28.3. Comparison 28 Azithromycin 3 to 5 days short-term treatment; Sensitivity analysis: funding not reported, Outcome 3 Treatment failure at 20 to 30 days.

Review: Short-course antibiotics for acute otitis media

Comparison: 28 Azithromycin 3 to 5 days short-term treatment; Sensitivity analysis: funding not reported

Outcome: 3 Treatment failure at 20 to 30 days




Analysis 29.1. Comparison 29 Azithromycin 3 to 5 days short-term treatment, adverse GI effects, Outcome 1 Gastrointestinal adverse effects.

Review: Short-course antibiotics for acute otitis media

Comparison: 29 Azithromycin 3 to 5 days short-term treatment, adverse GI effects

Outcome: 1 Gastrointestinal adverse effects




**Analysis 30.1. Comparison 30 Methods - combined control group versus twice counting tx, Outcome 1
Twice counting - short-acting antibiotic > 48 hours short-term treatment, treatment failure at 1 month or less.**

Review: Short-course antibiotics for acute otitis media

Comparison: 30 Methods - combined control group versus twice counting tx

Outcome: 1 Twice counting - short-acting antibiotic > 48 hours short-term treatment, treatment failure at 1 month or less




Analysis 30.2. Comparison 30 Methods - combined control group versus twice counting tx, Outcome 2 Combined control groups - short-acting antibiotic > 48 hours short-term treatment, treatment failure 1 month or less.

Review: Short-course antibiotics for acute otitis media

Comparison: 30 Methods - combined control group versus twice counting tx

Outcome: 2 Combined control groups - short-acting antibiotic > 48 hours short-term treatment, treatment failure 1 month or less




Analysis 30.3. Comparison 30 Methods - combined control group versus twice counting tx, Outcome 3 Twice counting - short-acting antibiotic > 48 hours short-term treatment, treatment failure at 1 to 19 days.

Review: Short-course antibiotics for acute otitis media

Comparison: 30 Methods - combined control group versus twice counting tx

Outcome: 3 Twice counting - short-acting antibiotic > 48 hours short-term treatment, treatment failure at 1 to 19 days




Analysis 30.4. Comparison 30 Methods - combined control group versus twice counting tx, Outcome 4 Combined control groups - short-acting antibiotic > 48 hours short-term treatment, treatment failure 1 to 19 days.

Review: Short-course antibiotics for acute otitis media

Comparison: 30 Methods - combined control group versus twice counting tx

Outcome: 4 Combined control groups - short-acting antibiotic > 48 hours short-term treatment, treatment failure 1 to 19 days




**Analysis 30.5. Comparison 30 Methods - combined control group versus twice counting tx, Outcome 5
Twice counting - short-acting antibiotic > 48 hours short-term treatment, treatment failure 20 to 30 days.**

Review: Short-course antibiotics for acute otitis media

Comparison: 30 Methods - combined control group versus twice counting tx

Outcome: 5 Twice counting - short-acting antibiotic > 48 hours short-term treatment, treatment failure 20 to 30 days




Analysis 30.6. Comparison 30 Methods - combined control group versus twice counting tx, Outcome 6 Combined control groups - short-acting antibiotic > 48 hours short-term treatment, treatment failure at 20 to 30 days.

Review: Short-course antibiotics for acute otitis media

Comparison: 30 Methods - combined control group versus twice counting tx

Outcome: 6 Combined control groups - short-acting antibiotic > 48 hours short-term treatment, treatment failure at 20 to 30 days




**Analysis 30.7. Comparison 30 Methods - combined control group versus twice counting tx, Outcome 7
Twice counting - short-acting antibiotic > 48 hours short-term treatment, treatment failure at 30 to 40 days.**

Review: Short-course antibiotics for acute otitis media

Comparison: 30 Methods - combined control group versus twice counting tx

Outcome: 7 Twice counting - short-acting antibiotic > 48 hours short-term treatment, treatment failure at 30 to 40 days




Analysis 30.8. Comparison 30 Methods - combined control group versus twice counting tx, Outcome 8 Combined control groups - short-acting antibiotic > 48 hours short-term treatment, treatment failure 30 to 40 days.

Review: Short-course antibiotics for acute otitis media

Comparison: 30 Methods - combined control group versus twice counting tx

Outcome: 8 Combined control groups - short-acting antibiotic > 48 hours short-term treatment, treatment failure 30 to 40 days




**Analysis 30.9. Comparison 30 Methods - combined control group versus twice counting tx, Outcome 9
Twice counting - short-acting antibiotic > 48 hours short-term treatment, treatment failure at 3 months or less.**

Review: Short-course antibiotics for acute otitis media

Comparison: 30 Methods - combined control group versus twice counting tx

Outcome: 9 Twice counting - short-acting antibiotic > 48 hours short-term treatment, treatment failure at 3 months or less




Analysis 30.10. Comparison 30 Methods - combined control group versus twice counting tx, Outcome 10 Combined control groups - short-acting antibiotic > 48 hours short-term treatment, treatment failure at 3 months or less.

Review: Short-course antibiotics for acute otitis media

Comparison: 30 Methods - combined control group versus twice counting tx

Outcome: 10 Combined control groups - short-acting antibiotic > 48 hours short-term treatment, treatment failure at 3 months or less



APPENDICES

Appendix I. Previous searches

The research librarian, in collaboration with the research team, developed and implemented search strategies designed to identify the highest quality of evidence for this review. The English and non-English language medical literature was searched using the MEDLINE and EMBASE databases to identify published clinical trials of AOM during the time period, January 1966 to July 1997. The Science Citation Index was utilised to identify additional trials which had cited relevant papers. The Current Contents/Life Sciences Index was searched for recently published AOM trials, not yet abstracted in MEDLINE.

Search updates were conducted with a modified search strategy using the following electronic resources for the years 1997 to August 2008 (exceptions stated in brackets) and no language restriction: MEDLINE, EMBASE, MEDLINE In-Process & Other Non-Indexed Citations, Cochrane Central Register of Controlled Trials (which contains the Cochrane Acute Respiratory Infections Group; this group handsearches journals pertinent to their content area and adds relevant trials to the registry), and International Pharmaceutical Abstracts, BIOSIS Previews (ISI Web of KnowledgeSM [3.0]), CINAHL Full Text (via EBSCOhost). The NLM (National Library of Medicine) Gateway, BioMed Central and OCLC PapersFirst and ProceedingsFirst were searched for identification of meeting abstracts. Additionally, trials registers such as Current Controlled Trials, ClinicalTrials.gov, the National Research Register, CRISP (Computer Retrieval of Information on Scientific Projects), the TRIP Database (Turning Research Into Practice), Scirus, Proquest Dissertations and Theses - Full Text (1861 to 2008), and Google Scholar were searched for additional unpublished controlled trials and reports. The

reference lists of relevant reviews and included studies were reviewed, and authors of included studies were contacted, as required (e.g. to clarify the source of population in cases of multiple publications or to seek additional data). For the search strategies, a combination of subject headings and keywords were adapted for each electronic resource using the following terms: 'otitis media', 'acute otitis media', 'ear infection', 'otorrhea', 'anti-infective agents', 'treatment outcome', 'antibacterial', 'antibiotic', 'bactericide' and 'antimicrobial'.

Appendix 2. Search strategies for 1997-2008 update

Electronic databases	Search strategies
MEDLINE OVID Version: rel10.5.2 1950 to August Week 2008 Searched: 1 August 2008 Results: 1892 Limits: Date: 1997 to 2008	1. exp Otitis Media/ 2. (acute adj5 (OM or otitis media or ear or infection?)).mp. 3. (OM or OME or AOM).ti,ab. 4. otorrh?ea.ti,ab. 5. or/1-4 6. exp Anti-Infective Agents/ 7. exp treatment outcome/ 8. (antibacter\$ or anti-bacter\$ or "anti bacter\$" or antibiotic\$ or anti-biotic\$ or "anti biotic\$" or bacteriocid\$ or antimicrob\$ or anti-microb\$ or "anti microb\$").mp. 9. or/6-8 10. clinical trial.pt. 11. randomized controlled trial.pt. 12. randomi?ed.ti,ab. 13. placebo.ti,ab. 14. dt.fs. 15. randomly.ti,ab. 16. trial.ti,ab. 17. groups.ti,ab. 18. or/10-17 19. animals/ 20. humans/ 21. 19 not (19 and 20) 22. 18 not 21 23. and/5,9,22 24. exp Infant/ 25. exp Child/ 26. Adolescent/ 27. Minors/ 28. exp Puberty/ 29. exp Pediatrics/ 30. infant\$.mp. 31. infancy.mp. 32. newborn\$.mp. 33. baby.mp. 34. babies.mp. 35. neonat\$.mp. 36. preterm\$.mp.

(Continued)

	37. prematur\$.mp. 38. postmatur\$.mp. 39. child\$.mp. 40. kid.mp. 41. kids.mp. 42. toddler\$.mp. 43. adolescen\$.mp. 44. teen\$.mp. 45. boy\$.mp. 46. girl.mp. 47. minor\$.mp. 48. pubert\$.mp. 49. pubescen\$.mp. 50. prepubescen\$.mp. 51. pediatric\$.mp. 52. paediatric\$.mp. 53. peadiatric\$.mp. 54. or/24-52 55. and/5,9,22,54 56. limit 55 to yr="1997 - 2007"
Ovid MEDLINE In-Process & Other Non-Indexed Citations OVID Version: rel10.5.2 1 August 2008 Searched: 1 August 2008 Results: 6 Limits: Date: 1997 to 2008	1. (acute adj5 (OM or otitis media or ear or infection?)).mp. 2. (OM or OME or AOM).ti,ab. 3. otorrh?ea.ti,ab. 4. or/1-3 5. (antibacter\$ or anti-bacter\$ or "anti bacter\$" or antibiotic\$ or anti-biotic\$ or "anti biotic\$" or bacteriocid\$ or antimicrob\$ or anti-microb\$ or "anti microb\$").mp. 6. (treatment adj5 outcome?).mp. [mp=title, original title, abstract, name of substance word, subject heading word] 7. or/5-6 8. "controlled clinical trial\$.mp. 9. "randomi?ed controlled trial\$.mp. 10. "research design\$.mp. 11. "clinical research".mp. 12. "random allocation".mp. 13. randomi?ed.ti,ab. 14. ("double blind" adj3 method\$.mp. 15. ("single blind" adj3 method\$.mp. 16. ((clin\$ or control\$) adj25 trial\$.mp. 17. ((singl\$ or doubl\$ or trebl\$ or tripl\$) adj25 (blind\$ or mask\$)).ti,ab. 18. placebo\$.ti,ab. 19. randomly.ti,ab. 20. or/8-19 21. infant\$.mp. 22. infancy.mp. 23. newborn\$.mp. 24. baby.mp.

(Continued)

	25. babies.mp. 26. neonat\$.mp. 27. preterm\$.mp. 28. prematur\$.mp. 29. postmatur\$.mp. 30. child\$.mp. 31. kid.mp. 32. kids.mp. 33. toddler\$.mp. 34. adolescen\$.mp. 35. teen\$.mp. 36. boy\$.mp. 37. girl.mp. 38. minor\$.mp. 39. pubert\$.mp. 40. pubescen\$.mp. 41. prepubescen\$.mp. 42. pediatric\$.mp. 43. paediatric\$.mp. 44. peadiatric\$.mp. 45. or/21-44 46. and/4,7,20,45 47. limit 46 to yr="1997 - 2007"
EMBASE OVID Version: rel10.5.2 1996 to 2008 Searched: 1 August 2008 Results: 2993 Limits: Date: 1997 to 2008	1. exp Otitis Media/ 2. (acute adj5 (OM or otitis media or ear or infection?)).mp. 3. (OM or OME or AOM).ti,ab. 4. otorrh?ea.ti,ab. 5. or/1-4 6. exp Antiinfective Agent/ 7. exp treatment outcome/ 8. (antibacter\$ or anti-bacter\$ or "anti bacter\$" or antibiotic\$ or anti-biotic\$ or "anti biotic\$" or bacteriocid\$ or antimicrob\$ or anti-microb\$ or "anti microb\$").mp. 9. or/6-8 10. exp clinical trial/ 11. randomi?ed.ti,ab. 12. placebo.ti,ab. 13. dt.fs. 14. randomly.ti,ab. 15. trial.ti,ab. 16. groups.ti,ab. 17. or/10-16 18. exp child/ 19. exp newborn/ 20. exp adolescent/ 21. exp puberty/ 22. exp pediatrics/ 23. infant\$.mp.

(Continued)

	24. infancy.mp. 25. newborn\$.mp. 26. baby.mp. 27. babies.mp. 28. neonat\$.mp. 29. preterm\$.mp. 30. prematur\$.mp. 31. postmatur\$.mp. 32. child\$.mp. 33. kid.mp. 34. kids.mp. 35. toddler\$.mp. 36. adolescen\$.mp. 37. teen\$.mp. 38. boy\$.mp. 39. girl.mp. 40. minor\$.mp. 41. pubert\$.mp. 42. pubescen\$.mp. 43. prepubescen\$.mp. 44. pediatric\$.mp. 45. paediatric\$.mp. 46. peadiatric\$.mp. 47. or/18-46 48. and/5,9,17,47 49. limit 48 to yr="1997 - 2007"
International Pharmaceutical Abstracts OVID Version: rel10.5.2 1970 to August 2008 Searched: 1 August 2008 Results: 45 Limits: Date: 1997 to 2008	1. (acute adj5 (OM or otitis media or ear or infection?)).mp. 2. (OM or OME or AOM).ti,ab. 3. otorrh?ea.ti,ab. 4. or/1-3 5. (antibacter\$ or anti-bacter\$ or "anti bacter\$" or antibiotic\$ or anti-biotic\$ or "anti biotic\$" or bacteriocid\$ or antimicrob\$ or anti-microb\$ or "anti microb\$").mp. 6. (treatment adj5 outcome?).mp. [mp=title, original title, abstract, name of substance word, subject heading word] 7. or/5-6 8. "controlled clinical trial\$.mp. 9. "randomi?ed controlled trial\$.mp. 10. "research design\$.mp. 11. "clinical research".mp. 12. "random allocation".mp. 13. randomi?ed.ti,ab. 14. ("double blind" adj3 method\$).mp. 15. ("single blind" adj3 method\$).mp. 16. ((clin\$ or control\$) adj25 trial\$).mp. 17. ((singl\$ or doubl\$ or trebl\$ or tripl\$) adj25 (blind\$ or mask\$)).ti,ab. 18. placebo\$.ti,ab.

(Continued)

	19. randomly.ti,ab. 20. or/8-19 21. infant\$.mp. 22. infancy.mp. 23. newborn\$.mp. 24. baby.mp. 25. babies.mp. 26. neonat\$.mp. 27. preterm\$.mp. 28. prematur\$.mp. 29. postmatur\$.mp. 30. child\$.mp. 31. kid.mp. 32. kids.mp. 33. toddler\$.mp. 34. adolescen\$.mp. 35. teen\$.mp. 36. boy\$.mp. 37. girl.mp. 38. minor\$.mp. 39. pubert\$.mp. 40. pubescen\$.mp. 41. prepubescen\$.mp. 42. pediatric\$.mp. 43. paediatric\$.mp. 44. peadiatric\$.mp. 45. or/21-44 46. and/4,7,20,45 47. limit 46 to yr="1997 - 2007"
EBM Reviews Cochrane Central Register of Controlled Trials 2 nd Quarter 2008 OVID Version: rel10.5.2 Searched: 1 August 2008 Results: 425 Limits: Date: 1997 to 2008	1. exp Otitis Media/ 2. (acute adj5 (OM or otitis media or ear or infection?)).mp. 3. (OM or OME or AOM).ti,ab. 4. otorrh?ea.ti,ab. 5. or/1-4 6. exp Anti-Infective Agents/ 7. exp treatment outcome/ 8. (antibacter\$ or anti-bacter\$ or "anti bacter\$" or antibiotic\$ or anti-biotic\$ or "anti biotic\$" or bacteriocid\$ or antimicrob\$ or anti-microb\$ or "anti microb\$").mp. 9. or/6-8 10. exp Infant/ 11. exp Child/ 12. Adolescent/ 13. Minors/ 14. exp Puberty/ 15. exp Pediatrics/ 16. infant\$.mp. 17. infancy.mp.

(Continued)

	18. newborn\$.mp. 19. baby.mp. 20. babies.mp. 21. neonat\$.mp. 22. preterm\$.mp. 23. prematur\$.mp. 24. postmatur\$.mp. 25. child\$.mp. 26. kid.mp. 27. kids.mp. 28. toddler\$.mp. 29. adolescen\$.mp. 30. teen\$.mp. 31. boy\$.mp. 32. girl.mp. 33. minor\$.mp. 34. pubert\$.mp. 35. pubescen\$.mp. 36. prepubescen\$.mp. 37. pediatric\$.mp. 38. paediatric\$.mp. 39. peadiatric\$.mp. 40. or/10-38 41. and/5,9,40 limit 41 to yr="1997 - 2007"
CINAHL Plus with Full Text (EBSCOhost) 1981-present Searched: 1 August 2008 Results: 532 Limits: Date: 1997 to 2008 Age Groups: Infant, Newborn 0-1 month, Infant, 1-23 months, Child, Preschool 2-5 years, Child, 6-12 years, Adolescence, 13-18 years	# Query S7(S5 and S1) S6(S5 and S1) S5(S4 or S3 or S2) S4(antibacter* or anti-bacter* or "anti bacter*" or antibiotic* or anti-biotic* or "anti biotic*" or bacteriocid* or antimicrob* or anti-microb* or "anti microb*") S3(MH "Treatment Outcomes+") S2(MH "Antiinfective Agents+") S1(MH "Otitis Media+") or (acute w5 OM OR acute w5 "otitis media" OR actue w5 ear OR acute w5 infection) or (ottorrh?ea or OM or OME or AOM)
BIOSIS Previews® (ISI Web of Knowledge SM [3.0]) 1969 to present Searched: 1 August 2008 Results: 1161 Limits: 1997 to 2008	Sets History #6 #5 AND #4 #5 TS=clinical trial* OR TS=research design OR TS=comparative stud* OR TS=evaluation stud* OR TS=controlled trial* OR TS= follow-up stud* OR TS=prospective stud* OR TS=random* OR TS=placebo* OR TS=(single blind*) OR TS=(double blind*) #4#3 OR #2 OR #1 #3 TS=ottorrhea

(Continued)

	#2 TS=(OM OR OME OR OME OR AOM) #1 TS="otitis media" OR TS=(acute SAME OM OR acute SAME "otitis media" OR acute SAME "ear infection")
OCLC PapersFirst (OCLC FirstSearch) Searched: 1 August 2008 Results: 25 + 3 Limits: 1997 to 2008	(kw: otitis w media) and (kw: antibiotic* OR kw: antibacter* OR kw: bacteriocid* OR kw: antimicrob*)
OCLC ProceedingsFirst (OCLC FirstSearch) Searched: 1 August 2008 Results: 1 Limits: 1997 to 2008	(kw: otitis w media) and (kw: antibiotic* OR kw: antibacter* OR kw: bacteriocid* OR kw: antimicrob*)
ProQuest® Dissertations & Theses Full Text 1861 to present Searched: 1 August 2008 Results: 62 = 3	((“otitis media” OR “ear infection” or otorrhea) AND (acute))

OVID databases:

RCT filter adapted from:

Cochrane Highly Sensitive Search Strategy (2005) Revision from

Glanville JM, Lefebvre C, Miles JNV, Camosso-Stefinovic J. How to identify randomized controlled trials in Medline: ten years on. J Med Libr Assoc 2006; 94(2):130-6

Appendix 3. Search Strategies 2009 Update

Database: **Ovid MEDLINE(R)** In-Process & Other Non-Indexed Citations and Ovid MEDLINE(R) <1950 to Present>

Search strategy:

1 otitis media.tw.

2 (OM or OME or AOM).tw.

3 (ear* adj3 (infect* or acute)).tw.

4 (otorrhea or otorrhoea).tw.

5 or/1-4

6 (antibacter* or anti-bacter* or anti bacter* or antibiotic* or anti-biotic* or anti biotic* or bacteriocid* or antimicrob* or anti-microb* or anti microb*).tw.

7 (amoxicillin* or amoxycillin* or penicillin* or cefprozil* or clarithromycin* or cefpodoxime* or cefaclor* or ceftriaxone* or azithromycin* or cefixime*).tw.

8 (treatment* adj5 outcome*).tw.

9 or/6-8

10 5 and 9

11 (infant* or infancy or newborn* or baby* or babies or neonat* or preterm* or prematur* or child* or kid or kids or toddler* or adolescen* or teen* or boy* or girl* or minor* or pubert* or pubescen* or prepubescen* or pediatric* or paediatric* or schoolchild* or school age* or preschool* or kindergar* or nursery school* or primary school* or secondary school* or elementary school* or high school* or highschool*).tw. (1599556)

12 10 and 11

13 (random* or placebo* or clinical trial* or research design* or doubl* blind* or singl* blind*).tw. (692296)

14 12 and 13

EMBASE.com

23. #18 AND #22
 22. #19 OR #20 OR #21
 21. ((singl* OR doubl*) NEAR/2 blind*):ab,ti
 20. random*:ab,ti OR placebo*:ab,ti OR factorial*:ab,ti OR crossover*:ab,ti OR 'cross-over':ab,ti OR 'cross over':ab,ti OR assign*:ab,ti OR allocat*:ab,ti OR volunteer*:ab,ti
 19. 'randomized controlled trial'/exp OR 'single blind procedure'/exp OR 'double blind procedure'/exp OR 'crossover procedure'/exp
 18. #12 AND #17
 17. #13 OR #14 OR #15 OR #16
 16. ((age* OR nursery OR primary OR secondary OR elementary OR high) NEAR/2 school*):ti,ab
 15. schoolchild*:ab,ti OR preschool*:ab,ti OR kindergar*:ab,ti OR highschool*:ab,ti
 14. infant*:ab,ti OR infancy:ab,ti OR newborn*:ab,ti OR baby*:ab,ti OR babies*:ab,ti OR neonat*:ab,ti OR preterm*:ab,ti OR prematur*:ab,ti OR postmatur*:ab,ti OR child*:ab,ti OR kid:ab,ti OR kids:ab,ti OR toddler*:ab,ti OR adolescen*:ab,ti OR teen*:ab,ti OR boy*:ab,ti OR girl*:ab,ti OR minor*:ab,ti OR pubert*:ab,ti OR pubescen*:ab,ti OR prepubescen*:ab,ti OR pediatric*:ab,ti OR paediatric*:ab,ti
 13. 'child'/exp OR 'newborn'/exp OR 'adolescent'/exp OR 'puberty'/exp OR 'pediatrics'/exp
 12. #6 AND #11
 11. #7 OR #8 OR #9 OR #10
 10. amoxicillin*:ab,ti OR amoxycillin*:ab,ti OR penicillin*:ab,ti OR cefprozil*:ab,ti OR clarithromycin*:ab,ti OR cefpodoxime*:ab,ti OR cefaclor*:ab,ti OR ceftriaxone*:ab,ti OR azithromycin*:ab,ti OR cefixime*:ab,ti
 9. antibacter*:ab,ti OR 'anti-bacterial':ab,ti OR 'anti-bacterials':ab,ti OR 'anti bacterial':ab,ti OR 'anti bacterials':ab,ti OR antibiotic*:ab,ti OR 'anti-biotic':ab,ti OR 'anti-biotics':ab,ti OR 'anti biotic':ab,ti OR bacteriocid*:ab,ti OR antimicrob*:ab,ti OR 'anti-microbial':ab,ti OR 'anti-microbials':ab,ti OR 'anti microbial':ab,ti OR 'anti microbials':ab,ti
 8. 'antiinfective agent'/exp
 7. 'treatment outcome'/exp
 6. #1 OR #2 OR #3 OR #4 OR #5
 5. otorrhoea:ab,ti OR otorrhea:ab,ti
 4. om:ab,ti OR ome:ab,ti OR aom:ab,ti
 3. ((infect* OR acute*) NEAR/5 ear*):ab,ti
 2. 'otitis media':ab,ti
 1. 'otitis media'/exp

FEEDBACK

'Short course antibiotics for acute otitis media' has severe errors of analysis and procedure and needs to be withdrawn, 29 November 2006

Summary

1. Data are counted twice as follows
 Study Short course Long course
 Boulesteix, 1995 11/124 11/118
 Cohen, 1997 26/186 31/184
 Hendrickse, 1988 14/74 6/77
 Hoberman, 1997a 57/197 24/178
 Hoberman, 1997b 57/197 40/189

Hoberman 1997a and b are the same study and so the short course data have been counted twice.

2. For both the Hoberman and Boulesteix studies the treatments being compared differ not only in terms of duration (short versus long) but also in terms of other aspects (for example, formulation, dose and dosing schedule). Thus as regards the comparison of short versus long they are biased.

Reply

1. Two studies ([Gooch 1996](#); [Hoberman 1997](#)) in the current review contained two arms comparing long course versus short course of antibiotics. In the original review the short arm was counted twice against each long arm of that study. In the update of this review, we have modified our methods and combined both long-course arms in a single comparison against the short-course arm, as recommended in the *Cochrane Handbook* ([Higgins 2008](#)).

The remaining articles ([Boulesteix 1995](#); [Cohen 1997](#); [Hendrickse 1988](#)) were checked to ensure that all patients enrolled in the study were accounted for only once. These three trials clearly indicate that each patient is only accounted for in one of the groups (short or long-course).

Hoberman 1997b has been removed from the updated review.

2. These minor differences were considered but were not deemed clinically important to the current question, and we don't believe this introduces any bias to the comparison.

Reply approved and added to review September 2009 by: Anita Kozyrskyj, Terry P Klassen, and Michael Moffatt

Contributors

Stephen Senn

Comment submitted 30 November 2006. Added to review 22 Decemer 2008.

WHAT'S NEW

Last assessed as up-to-date: 11 November 2009.

Date	Event	Description
19 June 2012	Review declared as stable	As of 19 June 2012, this Cochrane Review is no longer being updated, as there is high-quality evidence that treating children with acute otitis media with a short course (less than seven days) of antibiotics, compared to treatment with a long course (seven days or greater) of antibiotics, increases the likelihood of treatment failure in the short term, meaning further research is unlikely to change our confidence in the estimate of effect in our primary outcome. The review authors recommend that it is no longer necessary to update this review

HISTORY

Protocol first published: Issue 2, 1998

Review first published: Issue 2, 2000

Date	Event	Description
9 September 2010	Amended	Contact details updated.
18 November 2009	New citation required but conclusions have not changed	The searches were conducted in November 2009, with a total of 22 new trials added since the original review in 2000 (Adam 2000 ; Al-Ghamdi 1999 ; Arguedas 2005 ; Arrieta 2003 ; Block 2000 ; Block 2003 ; Block 2004 ; Catania 2004 ; Cohen 1998 ; Cohen 1999 ; Cohen 2000 ; Dagan 2000a ; Dagan 2000b ; de Jose 1998 ; Dunne 2003 ; Guvén 2006 ; Hoberman 2005 ; Kara 1998 ; Oguz 2003 ; Pessey 1999 ; Varsano 1997 ; Wang 2004). Two of the studies included in the first version of this review (Kozyrskyj 2000) were excluded due to the outcome not being reported as a treatment failure (Bain 1985 ; Jones 1986) and an additional three studies have been excluded; two were excluded due to use of cefibuten which has been deemed not to be clinically relevant (Roos 2000 ; Simon 1997); and one for the duration of antibiotic therapy being equivalent in both arms (Suzuki 2009). The 22 studies that were included in this updated review brought the 30-day findings closer to equivalence (primary outcome). This would suggest that any new studies would not change the current outcomes and the review authors recommend that it is no longer necessary to update this review
11 November 2009	New search has been performed	Searches conducted.
22 December 2008	Feedback has been incorporated	Feedback added.
12 March 2008	Amended	Converted to new review format.
30 January 2000	New search has been performed	Searches conducted.

CONTRIBUTIONS OF AUTHORS

Anita Kozyrskyj (AK), Terry Klassen (TK) and Michael Moffatt (MM) first conceived the review, presenting it as a meta-analysis in a journal ([Kozyrskyj 1998](#)). It was updated for *The Cochrane Library* with assistance from Krystal Harvey (KH) and Liza Bialy (LB).

DECLARATIONS OF INTEREST

None.

SOURCES OF SUPPORT

Internal sources

- Alberta Research Centre for Health Evidence, Canada.

External sources

- No sources of support supplied

INDEX TERMS

Medical Subject Headings (MeSH)

Acute Disease; Age Factors; Anti-Bacterial Agents [adverse effects; *therapeutic use]; Azithromycin [therapeutic use]; Ceftriaxone [adverse effects; therapeutic use]; Drug Administration Schedule; Otitis Media [*drug therapy]; Time Factors

MeSH check words

Child; Humans