

Childhood and Adolescent Sports-Related Overuse Injuries

KYLE J. CASSAS, M.D., and AMELIA CASSETTARI-WAYHS, M.D., *Methodist Health System, Dallas, Texas*

Youth sports participation carries an inherent risk of injury, including overuse injuries. Little leaguer's shoulder, a stress fracture of the proximal humerus that presents as lateral shoulder pain, usually is self-limited. Little leaguer's elbow is a medial stress injury; treatment consists of complete rest from throwing for four to six weeks followed by rehabilitation and a gradual throwing program. Spondylolysis is a stress fracture of the pars interarticularis. Diagnostic modalities include plain film radiography, bone scan, computed tomography, single photon emission computed tomography, and magnetic resonance imaging. Treatment usually is conservative. Spondylolisthesis is the forward or anterior displacement of one vertebral body over another and may be related to a history of spondylolysis. Diagnosis is made with plain film radiography and graded according to the amount of displacement. Osgood-Schlatter disease presents as anterior knee pain localized to the tibial tubercle. Diagnosis is made clinically, and most patients respond to conservative measures. Calcaneal apophysitis (or Sever's disease) is a common cause of heel pain in young athletes, presenting as pain in the posterior aspect of the calcaneus. (*Am Fam Physician* 2006;73:1014-22. Copyright © 2006 American Academy of Family Physicians.)

► **Patient information:** A handout on Osgood-Schlatter disease is available online at <http://familydoctor.org/135.xml>.

ACE This article exemplifies the AAFP 2006 Annual Clinical Focus on caring for children and adolescents.

Each year in the United States, approximately 30 million children and teenagers participate in organized sports.¹ Sports are the leading cause of injury in adolescents,² and the Centers for Disease Control and Prevention estimates that one half of all sports injuries in children are preventable^{2,3} with proper education and use of protective equipment. Children and adolescents may be particularly at risk for sports-related overuse injuries as a result of improper technique, poorly fitting protective equipment, training errors, and muscle weakness and imbalance. Most of these injuries can be managed conservatively with proper and timely diagnosis. Increasing numbers of chronic overuse injuries in young athletes may be related to limited recovery time from longer competitive seasons and year-round training.^{2,4}

The apophysis is a secondary center of ossification and a location for the insertion of a muscle tendon into bone.^{4,5} Overuse syndromes such as traction apophysitis may



ILLUSTRATION BY SCOTT BODELL

develop in young athletes when this growth center is unable to meet the demands placed on it during activity. Some sites of apophyseal injury include the knee (Osgood-Schlatter disease), heel (Sever's disease), and medial epicondyle (little leaguer's elbow).

Shoulder Pain in Overhead Sports

The demands of overhead sports (e.g., baseball, racquet sports, volleyball) put the shoulder at risk of injury. The etiology of shoulder pain may be varied and often is difficult to diagnose (*Table 1*). Rotator cuff syndrome is uncommon in children. However, conditions such as little leaguer's shoulder (i.e., gradual onset of pain in the proximal humerus during throwing⁶) are unique to overhead sports athletes with open growth plates. Little leaguer's shoulder is thought to be an overuse or stress injury of the proximal humeral physis and is not limited to baseball players; it also can be encountered in athletes involved in swimming, gymnastics, volleyball, and tennis.⁷ Parents and coaches must be aware that

SORT: KEY RECOMMENDATIONS FOR PRACTICE

<i>Clinical recommendation</i>	<i>Evidence rating</i>	<i>References</i>
Patients with little leaguer's shoulder should abstain from throwing for an average of three months.	C	6
Plain radiographs should be obtained when evaluating athletes with back pain of greater than three weeks' duration.	C	7,19
Single photon emission computed tomography should be used to diagnose acute spondylolysis, and computed tomography should be used to help stage and determine proper treatment.	C	20
Plain radiographs are rarely necessary for the diagnosis of Osgood-Schlatter disease or calcaneal apophysitis.	C	26,30

A = consistent, good-quality patient-oriented evidence; B = inconsistent or limited-quality patient-oriented evidence; C = consensus, disease-oriented evidence, usual practice, expert opinion, or case series. For information about the SORT evidence rating system, see page 956 or <http://www.aafp.org/afpsort.xml>.

any shoulder or elbow pain that persists in a young athlete may be a sign of an overuse or stress injury. Some evidence supports certain measures to reduce throwing and overuse injuries (i.e., adequate rest periods, icing after activity, evaluation for proper throwing mechanics, enforcement of proper pitch type and number for age, immediate reporting of pain during throwing or of persistent arm or shoulder pain).⁸⁻¹⁰

Little leaguer's shoulder should be suspected in young pitchers who throw breaking pitches (e.g., curveballs, sliders), which put increased demands on the shoulder and elbow.⁸ The average age of onset is 14 years; these patients typically present with lateral shoulder pain during throwing or overhead

activities.⁶ Most patients report insidious onset of symptoms that have been present for months and often delay seeking consultation until the pain increases or until there is a decrease in throwing velocity or control.⁷

Up to 70 percent of patients with little leaguer's shoulder have tenderness over the proximal and lateral portion of the humerus.^{6,7} Radiographs should be ordered to confirm the diagnosis: anteroposterior in internal and external rotation, lateral Y-view or axillary view, and comparison views of the uninvolved side. Classic findings include widening of the proximal humeral physis with or without physeal fragmentation, sclerosis, and demineralization (Figure 1).^{6,7}

Bone scan and magnetic resonance imaging (MRI) usually are unnecessary but may be considered if initial radiographs are negative and suspicion for the diagnosis is high, or if there is clinical suspicion for other pathology.

Treatment consists of relative rest from throwing for an average of three months, icing, and analgesic medications as needed for pain.⁶ Athletes are encouraged to maintain cardiovascular conditioning.⁶ Patients should begin strengthening exercises when they are comfortable and an interval throwing program when they are pain free. Evaluation of throwing mechanics also should be considered.

Elbow Pain in the Thrower

The elbow is the most common site of injury in young baseball players, and many of these injuries are preventable.^{4,11} Throwing

Little leaguer's shoulder should be suspected in young pitchers who throw breaking pitches such as curveballs and sliders.

TABLE 1

Causes of Shoulder Pain in Athletes

- Acromioclavicular sprain or injury
- Biceps tendonitis
- Bone tumor
- Brachial plexus injury
- Distal clavicle osteolysis
- Fracture
- Glenohumeral instability
- Impingement syndrome
- Labral pathology/SLAP tear
- Proximal humerus stress injury (little leaguer's shoulder)
- Referred pain
- Rotator cuff tendonitis or bursitis
- Thoracic outlet syndrome

SLAP = superior labrum anterior to posterior.

Childhood Sports Injuries

puts significant tension forces on the medial elbow and compression forces on the lateral radiocapitellar joint, leading to a spectrum of conditions that may include little leaguer's elbow (*Table 2*).^{9,12-14}

Little leaguer's elbow has been described as an apophysitis of the medial epicondyle in athletes between nine and 12 years of age.⁴ Most patients experience pain in the medial aspect of the elbow during throwing, and they may have decreased pitch velocity or control.^{9,15}

The diagnosis of little leaguer's elbow is made clinically and should be considered in throwers with medial elbow pain, even if symptoms are minimal.

The history should address hand dominance, position played, previous arm or elbow injuries, number and type of pitches per week, loss of control or velocity during throwing, decreased range of motion, numbness, and paresthesias. It also is important to ask about the number of

teams on which the athlete plays (club and/or school teams) and whether he or she has taken any time off from throwing during the year.

Examination may reveal swelling, loss of motion, and tenderness along the medial aspect of the elbow, distal arm, or proximal forearm.¹⁵ The diagnosis of little leaguer's elbow is made clinically and should be considered in throwers with medial elbow pain, even if symptoms are minimal. Initial evaluation includes elbow radiography with comparison views of the uninjured side.¹⁵

TABLE 2
Differential Diagnosis of Elbow Pain

Site	Possible diagnoses
Lateral	Avascular necrosis of the capitellum (Panner's disease; in children seven to 12 years of age)
	Osteochondritis dissecans (in children 12 to 16 years of age)
Medial	Flexor/pronator strain
	Medial apophysitis (little leaguer's elbow, in children nine to 12 years of age)
	Medial collateral ligament injury
	Medial epicondyle avulsion
Posterior	Ulnar neuritis
	Olecranon apophysitis
	Olecranon (posterior) impingement
	Olecranon osteochondrosis
Other	Triceps/olecranon tip avulsion
	Fracture
	Loose cartilaginous bodies
	Synovitis

Radiographs may be normal or may reveal hypertrophy of the medial epicondyle, bony fragmentation, apophyseal widening or avulsion (medial epicondyle), loose cartilaginous bodies, or osteochondral lesions.^{15,16}

MRI and nerve conduction studies may be useful for patients with normal radiographs and clinical suspicion for other conditions such as ulnar collateral ligament injury or radiculopathy.⁴




Figure 1. Comparison radiographs of little leaguer's shoulder in a 15-year-old baseball player. Note the widening of the proximal humeral epiphysis in the right shoulder (*A*) compared with the left shoulder (*B*).

TABLE 3

Possible Etiologies of Back Pain in Young Athletes

Ankylosing spondylitis
Congenital anomalies
Juvenile rheumatoid arthritis
Malignancy (e.g., leukemia, lymphoma)
Lumbar spondylolysis
Lumbar spondylolisthesis
Lumbosacral sprain
Osteochondrosis of the vertebrae (Scheuermann's disease)
Osteomyelitis
Scoliosis

Treatment consists of complete rest from throwing or pitching for at least four to six weeks; ice packs and analgesic medications may be used for swelling and pain. General conditioning, stretching, and core strengthening should be encouraged.^{4,9,15} A gradual and progressive (interval) throwing program may begin after the initial rest period. Most athletes are able to return to competitive pitching and throwing at 12 weeks.^{4,9,15} Orthopedic consultation is indicated for patients with loose cartilaginous bodies, avulsion fractures, and osteochondritis dissecans, and for patients whose symptoms do not improve with conservative treatment.¹²

To help prevent little leaguer's elbow and shoulder, the American Academy of Pediatrics recommends limiting the number of pitches to 200 per week or 90 pitches per outing.¹⁷ However, the USA Baseball Medical and Safety Advisory Committee recommends more conservative pitch counts (i.e., 75 to 125 pitches per week or 50 to 75 pitches per outing, depending on age).¹⁰ Other preventive measures include a pre-season conditioning program, instruction on proper pitching mechanics, and allowing time during the early part of the season to gradually increase the amount and intensity of throwing.¹⁷

Back Pain

Back pain in young athletes may be caused by many conditions (Table 3). However, primary care physicians should maintain a high index of suspicion for spondylolysis and spondylolisthesis because there is a higher incidence of these conditions in athletes.^{18,19}

Spondylolysis occurs in approximately 6 percent of the general population but may contribute to nearly 50 percent of cases of back pain in athletes.^{7,18,19} This vertebral defect (unilateral or bilateral) of the pars interarticularis is thought to be secondary to repetitive hyperextension of the posterior elements of the spine, leading to stress injury or fracture. Gymnastics, football, soccer, weightlifting, volleyball, ballet, and dance are known to put increased demands on the spine.^{7,18} Spondylolysis most commonly occurs at the fourth and fifth lumbar vertebral levels.^{7,18,19}

Most athletes with spondylolysis present with activity-related back pain that is exacerbated by hyperextension of the lumbar spine (e.g., lineman blocking, volleyball serving, gymnastic or cheerleading tumbling), progressing to pain during rest and daily activities. Examination may reveal a hyperlordotic posture, limited range of motion, and hamstring tightness with tenderness or pain in the affected region during single-leg hyperextension, commonly called the "stork" test (Figure 2).¹⁹

The initial diagnostic work-up for athletes




Figure 2. Single-leg hyperextension or "stork" test used to evaluate for pain associated with spondylolysis.

with back pain lasting longer than three weeks should include lumbar spine radiographs with anteroposterior, lateral, and bilateral oblique views (Figure 3²⁰).^{7,19} The classic appearance on the oblique view has been described as the “Scotty dog” with a “collar appearance” (Figure 4).⁷ Diagnosis of early spondylolysis with plain film radiography is difficult because the pars interarticularis may not have completely fractured, thus the “Scotty dog” will not be apparent.¹⁹

Some authors believe that plain film radiographs are insensitive and that further imaging often is necessary.^{18,20,21} Many studies¹⁸⁻²⁰ recommend the use of single photon emission computed tomography (SPECT) to help make an accurate diagnosis, assess the defect, and determine the potential for healing. One study²² reported SPECT imaging to have a 60 percent sensitivity and 85 percent specificity; however, the study population was not exclusive to young athletes. Whereas

Diagnosis and Treatment of Back Pain in Athletes



NOTE: Most athletes return to sports six months after diagnosis.

*—Consider bracing for athletes with continued symptoms after three weeks of rest. At three months, consider follow-up CT scan to assess for bony healing. Begin rehabilitation program at three months: spine stabilization and low-impact aerobic activity with progression as tolerated to sport-specific training.

†—Consider bracing for athletes with continued symptoms after three weeks of rest. Begin rehabilitation program once patient is asymptomatic: spine stabilization and low-impact aerobic activity with progression as tolerated to sport-specific training.

Figure 3. Suggested diagnostic and treatment algorithm for back pain in athletes. (SPECT = single photon emission computed tomography; CT = computed tomography.)

Adapted with permission from Standaert CJ. Spondylolysis in the adolescent athlete. *Clin J Sport Med* 2002;12:120.




ILLUSTRATION BY SCOTT BODELL

Figure 4. "Scotty dog" with a "collar appearance" seen on radiographs of patients with spondylolysis.

the SPECT scan is the most sensitive, computed tomography (CT) is highly specific and is useful after SPECT imaging for staging and treatment stratification.²⁰ MRI also may be useful for diagnostic evaluation, but there are limited data comparing CT, SPECT, and MRI.^{20,23,24}

Controversy exists regarding the ideal treatment of spondylolysis²⁰; however, accepted treatment includes relative rest from the offending activity, analgesic medications, physical therapy, and possibly bracing (for symptomatic patients after two to four weeks of rest). Bracing continues until the lesion is shown to be healed on radiographs or until the athlete is completely asymptomatic, which may take up to nine to 12 months.¹⁹ Some experts consider bracing to be a useful method of activity restriction rather than a means of immobilization for bony healing.²⁰ A high percentage of athletes with late-stage or terminal spondylolysis may not have bony healing, but rather an asymptomatic fibrous union.²⁰ If the condition is detected early and treated properly, the chances of healing with good long-term functional outcomes are improved.^{20,21} Physical therapy should focus on flexion activities, hamstring stretching, and core strengthening.

Without early detection and treatment, complete fracture and spondylolisthe-

sis with possible prolonged disability may occur. Most athletes respond to conservative management and return to full activity approximately six months after diagnosis²⁰; however, orthopedic consultation is warranted in recalcitrant cases. Athletes may return to sports once they have pain-free range of motion and normal strength and conditioning, and are able to perform sport-related skills without pain.²⁰

Spondylolisthesis is defined as the forward or anterior displacement of one vertebral body over another.²⁵ The etiology is unknown,²⁵ but the condition may be a result of trauma, overuse injuries, degeneration, or spondylolysis. Most patients with spondylolisthesis have a history of chronic back pain and present after an acute event or exacerbation of their pain. The examination may be similar to that for patients with spondylolysis and in severe cases may demonstrate a palpable step-off deformity of the lumbar spine.⁷

Diagnosis of spondylolisthesis is made with lateral lumbar plain film radiographs (Figure 5) and is graded according to the



Figure 5. Lateral radiograph of the lumbar spine showing grade 1 spondylolisthesis of the L5 vertebra on S1. There is a degree of sacral retrolisthesis. Note the pars defect (arrow).

amount of displacement (*Table 4*). Treatment for low-grade spondylolisthesis (i.e., up to 50 percent slippage) is similar to treatment for spondylolysis; however, to document stability and lack of further progression, patients should be followed with serial radiography at six-month intervals until they reach skeletal maturity.⁷ Patients with high-grade slippage (i.e., greater than 50 percent) and those who cannot be treated conservatively may need to be co-managed by an orthopedic or spine surgeon to guide treatment and assist in return-to-play decisions.⁷

Knee Pain

Knee pain is common in athletes (*Table 5*). **Osgood-Schlatter** disease often is encountered in children **10 to 15** years of age who participate in cutting and jumping sports such as soccer, basketball, gymnastics, and volleyball.²⁶ The exact cause of this condition is unknown, but it is thought to be secondary to repetitive microtrauma or a traction apophysitis of the tibial tuberosity.²⁶

Patients often have **anterior knee pain and swelling**, and 20 to 30 percent of patients report bilateral symptoms.²⁶⁻²⁸ Examination reveals tenderness and swelling at the tibial tubercle. **Radiographs** may reveal **fragmentation** and **irregular ossification** at the tibial tubercle but rarely are indicated unless there is suspicion of other injuries.²⁶ **Hip examination is important because some childhood hip conditions (e.g., slipped capital femoral epiphysis) can refer pain to the knee.**

Most patients respond to conservative treatment consisting of **rest from painful activities, icing, and analgesic medications** as needed for pain.²⁸ **Quadriceps stretching** as part of a strengthening program should be encouraged once symptoms are controlled. Corticosteroid injection of the tibial tubercle is not recommended because of potential complications such as subcutaneous atrophy.^{26,28}

Osgood-Schlatter disease usually is self-limited. However, some complications can

TABLE 4
Grading of Spondylolisthesis

Grade	Criterion
Grade I	Up to 25 percent slippage
Grade II	26 to 50 percent slippage
Grade III	51 to 75 percent slippage
Grade IV	76 to 100 percent slippage
Grade V	Greater than 100 percent slippage (spondyloptosis)

Information from reference 7.

TABLE 5
Differential Diagnosis of Anterior Knee Pain

- Fat pad hypertrophy/impingement (Hoffa's disease)
- Inferior patellar pole traction apophysitis (Sinding-Larsen disease)
- Patellofemoral pain syndrome
- Osgood-Schlatter disease
- Osteochondritis dissecans
- Patellar or quadriceps tendonitis
- Patellar stress fracture
- Prepatellar or infrapatellar bursitis
- Referred pain

Adapted with permission from Patel DR, Nelson TL. Sports injuries in adolescents. Med Clin North Am 2000;84:983-1007.

occur, such as mild discomfort with kneeling, a residual bony deformity (*Figure 6*), and painful ossicles in the distal patellar tendon.²⁶ Most patients are able to return to full activity within two to three weeks. The use of an infrapatellar strap for six to eight weeks may provide symptomatic relief during activity.²⁸

Heel Pain

Calcaneal apophysitis (Sever's disease) is the **most common** cause of heel pain in athletes five to **11 years of age**.²⁹ This **traction apophysitis** is secondary to repetitive microtrauma or **overuse** of the heel.^{14,27} Calcaneal

Spondylolysis and spondylolisthesis are common causes of back pain in young athletes.



Figure 6. Side (A) and frontal (B) views of a residual bony deformity associated with prior Osgood-Schlatter disease.



Figure 7. Typical location of tenderness associated with calcaneal apophysitis.

apophysitis is commonly seen in athletes participating in basketball, soccer, track, and other **running activities.**

Patients with calcaneal apophysitis may have activity-related pain in the posterior aspect of the heel, and **60 percent of patients report bilateral pain.**²⁷ Examination may reveal **tenderness on medial and lateral compression of the posterior calcaneus (Figure 7).** **Radiographs of the calcaneus may appear normal or may reveal a fragmented and sclerotic calcaneal apophysis.** Treatment consists of **activity modification, icing, stretching of the gastrocnemius-soleus complex, analgesic medications, and heel lifts or cushions.**³⁰ Most patients are able to return to pain-free activity within **three to six weeks.**

The Authors

KYLE J. CASSAS, M.D., is in private practice at Steadman Hawkins Clinic of the Carolinas, Simpsonville, S.C., and is program director for the Steadman Hawkins Primary Care Sports Medicine Fellowship, Greenville, S.C. He was assistant professor of family medicine at the University of Texas Southwestern Medical Center at Dallas and associate director of the sports medicine fellowship at Methodist Health System Family Medicine and Sports Medicine Center at the time the manuscript was written. He received his medical degree from the University of Nevada School of Medicine, Reno, and completed a surgical internship at Parkland Memorial Hospital in Dallas, Tex. Dr. Cassas completed a family medicine residency at St. Paul Medical Center in Dallas, where he was chief resident, and a sports medicine fellowship at Methodist Hospitals of Dallas.

AMELIA CASSETTARI-WAYHS, M.D., is assistant professor of family medicine at the University of Texas Southwestern Medical Center, Dallas, and assistant director of the Methodist Health System Family Medicine Residency Program and Sports Medicine Fellowship. She received her medical degree from Universidade Federal de Santa Catarina in Brazil and completed a family medicine residency at McAllen (Texas) Family Practice Residency Program.

Address correspondence to Kyle J. Cassas, M.D., Steadman Hawkins Clinic of the Carolinas, 727 SE Main St., Simpsonville, SC 29681 (e-mail: kcassas@ghs.org). Reprints are not available from the authors.

Author disclosure: Nothing to disclose.

REFERENCES

1. Adirim TA, Cheng TL. Overview of injuries in the young athlete. *Sports Med* 2003;33:75-81.
2. Emery CA. Risk factors for injury in child and adolescent sport: a systematic review of the literature. *Clin J Sport Med* 2003;13:256-68.

Childhood Sports Injuries

- U.S. Army Center for Health Promotion and Preventive Medicine. Youth sports injuries. Accessed online November 15, 2005, at: <https://chppm-www.apgea.army.mil/documents/YouthSportsInjuriesFS.pdf>.
- Benjamin HJ, Briner WW Jr. Little league elbow. *Clin J Sport Med* 2005;15:37-40.
- DeLee J, Drez D, Miller MD. *DeLee and Drez's Orthopaedic sports medicine: principles and practice*. 2nd ed. Philadelphia: Saunders, 2003.
- Carson WG Jr, Gasser SI. Little Leaguer's shoulder. A report of 23 cases. *Am J Sports Med* 1998;26:575-80.
- Patel DR, Nelson TL. Sports injuries in adolescents. *Med Clin North Am* 2000;84:983-1007.
- Lyman S, Fleisig GS, Andrews JR, Osinski ED. Effect of pitch type, pitch count, and pitching mechanics on risk of elbow and shoulder pain in youth baseball pitchers. *Am J Sports Med* 2002;30:463-8.
- Whiteside JA, Andrews JR, Fleisig GS. Elbow injuries in young baseball players. *Phys Sport Med* 1999;27:87-92,102.
- USA Baseball Medical and Safety Advisory Committee. Position statement on youth baseball injuries. Accessed online February 6, 2006, at: <http://www.usabaseball.com/ot/usab-med-safety-list.html>.
- Cain EL Jr, Dugas JR, Wolf RS, Andrews JR. Elbow injuries in throwing athletes: a current concepts review. *Am J Sports Med* 2003;31:621-35.
- Andrish JT. Osteochondritis dissecans in a young pitcher: why early recognition matters. *Phys Sport Med* 1997;25:85-90.
- Christopher NC, Congeni J. Overuse injuries in the pediatric athlete: evaluation, initial management, and strategies for prevention. *Clin Ped Emerg Med* 2002;3:118-28.
- Kaeding CK, Whitehead R. Musculoskeletal injuries in adolescents. *Prim Care* 1998;25:211-23.
- Behr CT, Altchek DW. The elbow. *Clin Sports Med* 1997;16:681-704.
- Hang DW, Chao CM, Hang YS. A clinical and roentgenographic study of little league elbow. *Am J Sports Med* 2004;32:79-84.
- Committee on Sports Medicine and Fitness. American Academy of Pediatrics: risk of injury from baseball and softball in children. *Pediatrics* 2001;107:782-4.
- Congen J, McCulloch J, Swanson K. Lumbar spondylolysis. A study of natural progression in athletes. *Am J Sports Med* 1997;25:248-53.
- Solomon R, Brown T, Gerbino PG, Micheli LJ. The young dancer. *Clin Sports Med* 2000;19:717-39.
- Standaert CJ. Spondylolysis in the adolescent athlete. *Clin J Sport Med* 2002;12:119-22.
- Miller SF, Congeni J, Swanson K. Long-term functional and anatomical follow-up of early detected spondylolysis in young athletes. *Am J Sports Med* 2004;32:928-33.
- Gregory PL, Batt ME, Kerslake RW, Scammell BE, Webb JF. The value of combining single photon emission computerised tomography and computerised tomography in the investigation of spondylolysis. *Eur Spine J* 2004;13:503-9.
- Campbell RS, Grainger AJ, Hide IG, Papastefanou S, Greenough CG. Juvenile spondylolysis: a comparative analysis of CT, SPECT and MRI. *Skeletal Radiol* 2005;34:63-73.
- Cohen E, Stuecker RD. Magnetic resonance imaging in diagnosis and follow-up of impending spondylolysis in children and adolescents: early treatment may prevent pars defects. *J Pediatr Orthop B* 2005;14:63-7.
- Ikata T, Miyake R, Katoh S, Morita T, Murase M. Pathogenesis of sports-related spondylolisthesis in adolescents. Radiographic and magnetic resonance imaging study. *Am J Sports Med* 1996;24:94-8.
- Wall EJ. Osgood-Schlatter disease: practical treatment for a self-limited condition. *Phys Sport Med* 1998;26:29-34.
- Marx JA, Hockberger RS, Walls RM, Adams J. *Rosen's Emergency medicine: concepts and clinical practice*. 5th ed. St. Louis: Mosby, 2002.
- Bloom OJ, Mackler L, Barbee J. Clinical inquiries. What is the best treatment for Osgood-Schlatter disease? *J Fam Pract* 2004;53:153-6.
- Schroeder BM. American College of Foot and Ankle Surgeons: diagnosis and treatment of heel pain. *Am Fam Physician* 2002;65:1686,1688.
- Micheli LJ, Ireland ML. Prevention and management of calcaneal apophysitis in children: an overuse syndrome. *J Pediatr Orthop* 1987;7:34-8.